

ORBITS OF DISCRETE SUBGROUPS ON A SYMMETRIC SPACE AND THE FURSTENBERG BOUNDARY

ALEXANDER GORODNIK AND HEE OH

Dedicated to Hillel Furstenberg with respect and admiration

ABSTRACT. Let X be a symmetric space of noncompact type and Γ a lattice in the isometry group of X . We study the distribution of orbits of Γ acting on the symmetric space X and its geometric boundary $X(\infty)$ generalizing the main equidistribution result of Margulis' thesis to higher rank symmetric spaces. More precisely, for any $y \in X$ and $b \in X(\infty)$, we investigate the distribution of the set $\{(y\gamma, b\gamma^{-1}) : \gamma \in \Gamma\}$ in $X \times X(\infty)$. It is proved, in particular, that the orbits of Γ in the Furstenberg boundary are equidistributed, and that the orbits of Γ in X are equidistributed in "sectors" defined with respect to a Cartan decomposition. Our main tools are the strong wavefront lemma and the equidistribution of solvable flows on homogeneous spaces, which we obtain using Shah's result based on Ratner's measure classification theorem.

CONTENTS

1. Introduction	1
2. Main ingredients of the proofs	7
3. Cartan decomposition and the strong wavefront lemma	10
4. Contractions on G/B	15
5. Volume estimates	18
6. Equidistribution of solvable flows	21
7. Distribution of lattice points in sectors and the boundary	28
8. Distribution of lattice points in bisectors	32
9. Measure-preserving lattice actions	35
References	36

1. INTRODUCTION

Let \mathbb{D} denote the hyperbolic unit disc and Γ a torsion free discrete subgroup of the isometry group of \mathbb{D} such that \mathbb{D}/Γ has finite area. The geometric boundary of \mathbb{D} is the space of the equivalence classes of geodesic rays in \mathbb{D} . It can be

The first author is partially supported by NSF grant 0400631.

The second author partially supported by NSF grant 0333397.

identified with the unit circle \mathbb{S} . Note that the action of Γ on \mathbb{D} extends to the geometric boundary of \mathbb{D} .

Let $x \in \mathbb{D}$. We denote by $B_T(x)$ the ball of radius T centered at x . For an arc $\Omega \subset \mathbb{S}$, the sector $\mathcal{S}_x(\Omega)$ in \mathbb{D} is defined to be the set of points $z \in \mathbb{D}$ such that the end point of the geodesic ray from x to z lies in Ω . Denote by m_x the unique probability measure on \mathbb{S} invariant under the isometries that fix the point x . Then:

A. For any $x, y \in \mathbb{D}$, $b \in \mathbb{S}$ and an arc $\Omega \subset \mathbb{S}$,

$$\#\{\gamma \in \Gamma : b\gamma^{-1} \in \Omega, y\gamma \in B_T(x)\} \sim_{T \rightarrow \infty} m_y(\Omega) \cdot \frac{\text{Area}(B_T(x))}{\text{Area}(\mathbb{D}/\Gamma)}.$$

B. For any $x, y \in \mathbb{D}$ and an arc $\Omega \subset \mathbb{S}$,

$$\#\{\gamma \in \Gamma : y\gamma \in \mathcal{S}_x(\Omega) \cap B_T(x)\} \sim_{T \rightarrow \infty} m_x(\Omega) \cdot \frac{\text{Area}(B_T(x))}{\text{Area}(\mathbb{D}/\Gamma)}.$$

C. For every $x, y \in \mathbb{D}$, $b \in \mathbb{S}$ and arcs $\Omega_1, \Omega_2 \subset \mathbb{S}$,

$$\begin{aligned} \#\{\gamma \in \Gamma : y\gamma \in \mathcal{S}_x(\Omega_1) \cap B_T(x), b\gamma^{-1} \in \Omega_2\} \\ \sim_{T \rightarrow \infty} m_x(\Omega_1)m_y(\Omega_2) \cdot \frac{\text{Area}(B_T(x))}{\text{Area}(\mathbb{D}/\Gamma)}. \end{aligned}$$

Statement (A) may be deduced from the work of A. Good [G]. Statement (B) was shown by P. Nicholls [N] (see also [Sh]). Statement (C), which was proved by G. Margulis [M] in his thesis for cocompact Γ , shows that the equidistribution phenomena in (A) and (B) are indeed independent.




FIGURE 1

The main purpose of this paper is to obtain an analog of statement (C) (note that (C) implies both (A) and (B)) for an arbitrary Riemannian symmetric space of noncompact type (see Theorems 1.1 and 1.2 below). We also generalize statement (B) to the equidistribution of lattice points in a connected noncompact semisimple Lie group G with finite center with respect to both K -components in a Cartan decomposition $G = KA^+K$ (see Theorem 1.6 below).

Let X be a Riemannian symmetric space of noncompact type and $X(\infty)$ the geometric boundary of X (that is, the space of equivalence classes of geodesic rays in X). Denote by G the identity component of the isometry group of X acting on X from the right. Let Γ be a lattice in G , i.e., a discrete subgroup with finite covolume. The action of G on X extends to $X(\infty)$.

For $x \in X$, we denote by $B_T(x)$ the Riemannian ball of radius T centered at x , by K_x the stabilizer of x in G , and by ν_x the probability Haar measure on K_x . For $x \in X$ and $b \in X(\infty)$, we denote by $m_{b,x}$ the unique probability K_x -invariant measure supported on the orbit $bG \subset X(\infty)$ (note that G acts transitively on $X(\infty)$ only when the rank of X is one, and that K_x acts transitively on each G -orbit in $X(\infty)$). Fix a closed Weyl chamber $\mathcal{W}_x \subset X$ at x . According to the Cartan decomposition, we have $X = \mathcal{W}_x K_x$. Let M_x denote the stabilizer of \mathcal{W}_x in K_x .

The following is one of our main theorems:

Theorem 1.1. *For $x, y \in X$, $b \in X(\infty)$, and any Borel subsets $\Omega_1 \subset K_x$ and $\Omega_2 \subset bG$ with boundaries of measure zero,*

$$\#\{\gamma \in \Gamma : y\gamma \in \mathcal{W}_x \Omega_1 \cap B_T(x), b\gamma^{-1} \in \Omega_2\} \sim_{T \rightarrow \infty} \nu_x(M_x \Omega_1) m_{b,y}(\Omega_2) \cdot \frac{\text{Vol}(B_T)}{\text{Vol}(G/\Gamma)},$$

where $\text{Vol}(B_T)$ denotes the volume of a ball of radius T in X .

We deduce Theorem 1.1 from a stronger result on the level of Lie groups. Fix the following data:

- G – a connected noncompact semisimple Lie group with finite center,
- G_n – the product of all noncompact simple factors of G ,
- $G = K_1 A^+ K_1$ – a Cartan decomposition of G ,
- d – an invariant metric on $K_1 \backslash G$,
- K_2 – a maximal compact subgroup of G ,
- Q – a closed subgroup of G that contains a maximal connected split solvable subgroup.

Recall that a solvable subgroup S is called *split* if the eigenvalues of any element of $\text{Ad}(S)$ are real for the adjoint representation $\text{Ad} : G \rightarrow \text{GL}(\text{Lie}(G))$. It is well-known that a maximal connected split solvable subgroup is a subgroup of the form AN for an Iwasawa decomposition $G = KAN$. Thus, $G = K_2 Q$. Denote by ν_1 and ν_2 the probability Haar measure on K_1 and K_2 respectively.

Let M_1 be the centralizer of A in K_1 and $M_2 = K_2 \cap Q$. Since any two maximal compact subgroups of G are conjugate to each other, there exists $g \in G$ such that $K_2 = g^{-1}K_1g$. Let Γ be a lattice in G such that $\overline{\Gamma G_n Q} = G$.

Theorem 1.2. *For any Borel subsets $\Omega_1 \subset K_1$ and $\Omega_2 \subset K_2$ with boundaries of measure zero,*

$$\begin{aligned} \#\{\gamma \in \Gamma \cap g^{-1}K_1A^+\Omega_1 \cap \Omega_2Q : d(K_1, K_1g\gamma) < T\} \\ \sim_{T \rightarrow \infty} \nu_1(M_1\Omega_1)\nu_2(\Omega_2M_2) \cdot \frac{\text{Vol}(G_T)}{\text{Vol}(G/\Gamma)}, \end{aligned}$$

where $\text{Vol}(G_T)$ denotes the volume of a Riemannian ball of radius T in G .

To understand the presence of M_1 and M_2 in the above asymptotics, observe that $K_1A^+\Omega_1 = K_1A^+M_1\Omega_1$ and $\Omega_2M_2Q = \Omega_2Q$.

Remark 1.3. We mention that the continuous version of Theorem 1.2 does not seem obvious either. The method of the proof of Theorem 1.2 also yields the following volume asymptotics:

$$\begin{aligned} \text{Vol}(\{h \in g^{-1}K_1A^+\Omega_1 \cap \Omega_2Q : d(K_1, K_1gh) < T\}) \\ \sim_{T \rightarrow \infty} \nu_1(M_1\Omega_1)\nu_2(\Omega_2M_2) \cdot \frac{\text{Vol}(G_T)}{\text{Vol}(G/\Gamma)}. \end{aligned}$$

Remark 1.4. If in Theorem 1.2 we replace $K_1A^+\Omega_1$ by $\Omega_1A^+K_1$, then the statement of the theorem is false. In fact, we can show that there exist nonempty open subsets $\Omega_1 \subset K_1$ and $\Omega_2 \subset K_2$ such that

$$\lim_{T \rightarrow \infty} \frac{1}{\text{Vol}(G_T)} \text{Vol}(\{h \in g^{-1}\Omega_1A^+K_1 \cap \Omega_2Q : d(K_1, K_1gh) < T\}) = 0.$$

To state yet another generalization of statement (B), we fix a Cartan decomposition $G = KA^+K$ and an invariant Riemannian metric d on $K \backslash G$. Let Γ be any lattice in G . Recall that it was shown in [DRS] and [EM] that for a lattice Γ in G ,

$$(1.5) \quad \#\{\gamma \in \Gamma : d(K, Kg\gamma) < T\} = \#(\Gamma \cap g^{-1}KA_T^+K) \sim_{T \rightarrow \infty} \frac{\text{Vol}(G_T)}{\text{Vol}(G/\Gamma)},$$

where $A_T^+ = \{a \in A^+ : d(K, Ka) < T\}$. The following theorem is a generalization of this result:

Theorem 1.6. *For $g \in G$ and any Borel subsets $\Omega_1 \subset K$ and $\Omega_2 \subset K$ with boundaries of measure zero,*

$$\#(\Gamma \cap g^{-1}\Omega_1A_T^+M\Omega_2) \sim_{T \rightarrow \infty} \frac{\text{Vol}(g^{-1}\Omega_1A_T^+M\Omega_2)}{\text{Vol}(G/\Gamma)} = \nu(\Omega_1M)\nu(M\Omega_2) \cdot \frac{\text{Vol}(G_T)}{\text{Vol}(G/\Gamma)},$$

where M is the centralizer of A^+ in K , and ν is the probability Haar measure on K .

We now present several corollaries of (the methods of) the above theorems.

1.1. Lattice action on the Furstenberg boundary. For a connected semisimple Lie group G with finite center, the Furstenberg boundary of G is identified with the quotient space G/P where P is a minimal parabolic subgroup of G (see [GJT, Ch. IV]). In the rank one case, the Furstenberg boundary G/P coincides with the geometric boundary $X(\infty)$ of the symmetric space X of G . In the higher rank case, G/P is isomorphic to the G -orbit in $X(\infty)$ of any regular geodesic class and can be identified with the space of asymptotic classes of Weyl chambers in X .

It is well-known that the action of a lattice Γ on G/P is minimal, i.e., every Γ -orbit is dense ([Mo, Lemma 8.5]). A natural question is whether each Γ -orbit in G/P is equidistributed. Corollary 1.7, which is a special case of Theorem 1.2, implies an affirmative answer in a much more general setting.

Let d denote an invariant Riemannian metric on the symmetric space $X \simeq K \backslash G$, where K is a maximal compact subgroup of G .

Corollary 1.7. *Let Q be a closed subgroup of G containing a maximal connected split solvable subgroup of G and $g \in G$. Denote by ν_g the unique $g^{-1}Kg$ -invariant probability measure on G/Q . Let $b \in G/Q$ and Γ be a lattice in G such that $\overline{\Gamma G_n b} = G/Q$. Then for any Borel subset $\Omega \subset G/Q$ such that $\nu_g(\partial\Omega) = 0$,*

$$\#\{\gamma \in \Gamma : \gamma b \in \Omega, d(K, Kg\gamma) < T\} \sim_{T \rightarrow \infty} \nu_g(\Omega) \cdot \frac{\text{Vol}(G_T)}{\text{Vol}(G/\Gamma)}.$$

It follows from the result of N. Shah [S, Theorem 1.1] and Ratner's topological rigidity [R2] that the condition $\overline{\Gamma G_n b} = G/Q$ is equivalent to the density of the orbit Γb in G/Q .

In the case when Q is a parabolic subgroup of G , a different proof of this result, which is based on ideas developed in [Ma], is recently given in [GM].

In the last decade or so there have been intensive studies on the equidistribution properties of lattice points on homogeneous spaces of G using various methods from analytic number theory, harmonic analysis and ergodic theory ([DRS], [EM],[EMM], [EMS], [GO], [EO], [Go], [L], [Ma], [No], etc.). Of particular interest is the case when the homogeneous space is a real algebraic variety. While most of the attention in this direction is focused on the case of affine homogeneous varieties, there is not so much work done for the projective homogeneous varieties, except for recent works [Go] and [Ma]. In [Go], one studies the distribution of lattice orbits on the real projective homogeneous varieties of $G = \text{SL}_n(\mathbb{R})$ with respect to the norm given by $\|g\| = \sqrt{\sum g_{ij}^2}$, $g \in \text{SL}_n(\mathbb{R})$. In [Ma], one investigates the distribution of lattice orbits on the boundary of a real hyperbolic

space. Corollary 1.7 extends both results by proving that an orbit of a lattice in a connected noncompact semisimple real algebraic group G is equidistributed on any projective homogeneous variety of G (with respect to a Riemannian metric).

More generally, we state the following conjecture:

Conjecture 1.8. *Let G be a connected semisimple Lie group with finite center, Γ a lattice in G , and Y a compact homogeneous space of G . Then every dense orbit of Γ in Y is equidistributed, i.e., there exists a smooth measure ν on Y such that for any $y \in Y$ with $\overline{\Gamma y} = Y$ and for any Borel set $\Omega \subset Y$ with boundary of measure zero,*

$$\#\{\gamma \in \Gamma : \gamma y \in \Omega, d(K, K\gamma) < T\} \sim_{T \rightarrow \infty} \nu(\Omega) \cdot \frac{\text{Vol}(G_T)}{\text{Vol}(G/\Gamma)}.$$

The structure of compact homogeneous spaces of G was studied in [Mor]. We note that the case of the conjecture when $Y = G/\Gamma$ for a cocompact lattice is also known (see Theorem 1.9).

1.2. Measure-preserving lattice actions. Let G be a connected semisimple noncompact Lie group with finite center and Γ_1, Γ_2 lattices in G . We consider the action of Γ_1 on G/Γ_2 . Let d be an invariant Riemannian metric on the symmetric space $K \backslash G$.

Theorem 1.9. *Suppose that for $y \in G/\Gamma_2$, the orbit $\Gamma_1 y$ is dense in G/Γ_2 . Then for any $g \in G$ and any Borel subset $\Omega \subset G/\Gamma_2$ with boundary of measure zero,*

$$\{\gamma \in \Gamma_1 : \gamma y \in \Omega, d(K, Kg\gamma) < T\} \sim_{T \rightarrow \infty} \frac{\text{Vol}(\Omega) \cdot \text{Vol}(G_T)}{\text{Vol}(G/\Gamma_1) \text{Vol}(G/\Gamma_2)},$$

where all volumes are computed with respect to one fixed Haar measure on G .

For example, Theorem 1.9 applies to the case when G is a simple connected noncompact Lie group, and Γ_1 and Γ_2 are noncommensurable lattices in G . (Recall that the lattices are called *commensurable* if $\Gamma_1 \cap \Gamma_2$ has finite index in both Γ_1 and Γ_2 .) It was first observed by Vatsal (see [V]) that $\Gamma_1 \Gamma_2$ is dense in G/Γ_2 . This is a (simple) consequence of Ratner's topological rigidity [R2].

Theorem 1.9 was proved in [O] for $G = \text{SL}_n(\mathbb{R})$ equipped with the norm $\|g\| = \sqrt{\sum g_{ij}^2}$ and in [GW] for general semisimple Lie groups without compact factors.

1.3. Acknowledgments. The first author wishes to thank for the hospitality of the Department of Mathematics in California Institute of Technology during August 2003 when most of this work has been done.

2. MAIN INGREDIENTS OF THE PROOFS

2.1. The strong wavefront lemma. The following theorem is a basic tool which enables us to reduce the counting problems for Γ as in Theorems 1.2 and 1.6 to the study of continuous flows on the homogeneous space $\Gamma \backslash G$.

Let G be a connected noncompact semisimple Lie group with finite center, $G = KA^+K$ a Cartan decomposition, and M the centralizer of A in K .

Theorem 2.1 (The strong wavefront lemma). *Let \mathcal{C} be any closed subset of A^+ with a positive distance from the walls of A^+ . Then for any neighborhoods U_1, U_2 of e in K and V of e in A , there exists a neighborhood \mathcal{O} of e in G such that for any $g = k_1 a k_2 \in KCK$,*

- (1) $g\mathcal{O} \subset (k_1 U_1)(aVM)(k_2 U_2)$;
- (2) $\mathcal{O}g \subset (k_1 U_1)(aVM)(k_2 U_2)$.

The strong wavefront lemma has been recently generalized to affine symmetric spaces in [GOS].

Remark 2.2. One can check that Theorem 2.1 fails if the set \mathcal{C} contains a sequence that converges to a point in a wall of the Weyl chamber A^+ .

Theorem 2.1 has several geometric implications for the symmetric space $K \backslash G$:

- *Strengthening of the wavefront lemma.* Recall that the wavefront lemma introduced by Eskin and McMullen in [EM] says that for any neighborhood \mathcal{O}' of e in G , there exists a neighborhood \mathcal{O} of e in G such that

$$(2.3) \quad a\mathcal{O} \subset \mathcal{O}'aK \quad \text{for all } a \in A^+.$$

To see that our strong wavefront lemma (1) implies the wavefront lemma for $a \in A^+$ with at least a fixed positive distance from the walls of A^+ , note that \mathcal{O}' contains $U_1 V$ for some neighborhood U_1 of e in K and some neighborhood V of e in A , and hence

$$U_1 a V M K = U_1 V a K \subset \mathcal{O}' a K$$

By Theorem 2.1 (1), there exists a neighborhood \mathcal{O} of e such that

$$a\mathcal{O} \subset U_1 a V M K.$$

Thus, $a\mathcal{O} \subset \mathcal{O}' a K$.

To illustrate geometric meaning of the strong wavefront lemma (1), we consider the unit disc \mathbb{D} equipped with the standard hyperbolic metric and the geodesic flow g_t on the unit tangent bundle $T^1(\mathbb{D})$ that transports a vector distance t along the geodesic to which it is tangent. Note that with the identification $T^1(\mathbb{D}) \simeq \mathrm{PSL}_2(\mathbb{R})$, the geodesic flow g_t corresponds to the left multiplication by $\begin{pmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{pmatrix}$. Let $p \in \mathbb{D}$ and let $K \subset T^1(\mathbb{D})$ be the preimage of p under the projection map $\pi : T^1(\mathbb{D}) \rightarrow \mathbb{D}$. Note that




FIGURE 2

K consists of vectors lying over p and pointing in all possible directions, and $g_t(K)$ consists of the unit vectors normal to the sphere $S_t(p) \subset \mathbb{D}$ of radius t . The wavefront lemma (see (2.3)) implies that one can find a neighborhood $\mathcal{O} \subset T^1(\mathbb{D})$ of a vector v based at p such that $g_t(\mathcal{O})$ remains close to $g_t(K)$ uniformly for every $t \geq 0$.

However, this does not compare $g_t(\mathcal{O})$ with the vector $g_t(v)$, but rather with the set $g_t(K)$. Theorem 2.1 (1) says that we may choose a neighborhood \mathcal{O} of v in $T^1(\mathbb{D})$ so that $g_t(\mathcal{O})$ is close to the vector $g_t(v)$ uniformly on t in both angular and radial components (see Fig. 2).

- *Uniform openness of the map $K \times A^+ \times K \rightarrow G$.* The product map

$$K \times (\text{interior of } A^+) \times K \rightarrow G$$

is a diffeomorphism onto a dense open subset in G , and in particular, it is an open map. Theorem 2.1 (2) shows that this map is *uniformly open* with respect to the base of neighborhoods $\mathcal{O}g$, where \mathcal{O} is a neighborhood of e in G and $g \in G$, on any subset contained in $K \times A^+ \times K$ with a positive distance from the walls of A^+ .

We illustrate the geometric meaning of this property for the case of the hyperbolic unit disc \mathbb{D} . It follows from Theorem 2.1 (2) that for every neighborhood \mathcal{O}' of e in G , there exists a neighborhood \mathcal{O} of e in G such that for every $a \in A^+$ with at least a fixed positive distance from the walls of A^+ ,

$$\mathcal{O}a \subset Ka\mathcal{O}'.$$

This implies that for every open subset $\mathcal{O}' \subset \mathbb{T}^1(\mathbb{D})$, $p \in \mathcal{O}'$, and any $t > 0$, the set $\pi(g_t(\mathcal{O}'))$ contains a ball centered at $\pi(g_t(p))$ with a radius independent of t .

- *Well-roundness of bisectors.* Theorem 2.1 (2) also implies the following corollary, which is a generalization of the well-known property that the Riemannian balls are well-rounded (a terminology used in [EM]). Note that the Riemannian ball $\{g \in G : d(K, Kg) < T\}$ is of the form KA_T^+K , where $A_T^+ = \{a \in A^+ : d(K, Ka) < T\}$.

Corollary 2.4. *For Borel subsets $\Omega_1, \Omega_2 \subset K$ whose boundary has measure zero, the family $\{\Omega_1 A_T^+ \Omega_2 : T > 0\}$ of bisectors is well-rounded, i.e., for every $\varepsilon > 0$, there exists a neighborhood \mathcal{O} of e in G such that*

$$\text{Vol}(\mathcal{O} \cdot \partial(\Omega_1 A_T^+ \Omega_2)) \leq \varepsilon \cdot \text{Vol}(\Omega_1 A_T^+ \Omega_2)$$

for all $T > 0$.

2.2. Uniform distribution of solvable flows. Using the strong wavefront lemma, Theorem 1.2 is deduced from Theorem 2.5 below, which is also of independent interest from the viewpoint of ergodic theory.

Let K be a maximal compact subgroup of G with the probability Haar measure ν . Let Q be a closed subgroup of G containing a maximal connected split solvable subgroup of G , and let ρ be a right invariant Haar measure on Q . Fix a Cartan decomposition $G = KA^+K$ and $g \in G$. For $T > 0$ and a subset $\Omega \subset K$, we define

$$Q_T(g, \Omega) = \{q \in Q : q \in g^{-1}KA^+\Omega, d(K, Kq) < T\}.$$

If $\Omega = K$ and $g = e$, the set $Q_T(g, \Omega)$ is simply $\{q \in Q : d(K, Kq) < T\}$. Recall that G_n denotes the product of all noncompact simple factors of G .

Theorem 2.5. *Let G be realized as a closed subgroup of a Lie group L . Let Λ be a lattice in L . Suppose that for $y \in \Lambda \backslash L$, the orbit yG_n is dense in $\Lambda \backslash L$. Then for any Borel subset $\Omega \subset K$ with boundary of measure zero and $f \in C_c(\Lambda \backslash L)$,*

$$\lim_{T \rightarrow \infty} \frac{1}{\rho(Q_T(g, K))} \int_{Q_T(g, \Omega)} f(yq^{-1}) d\rho(q) = \frac{\nu(M\Omega)}{\mu(\Lambda \backslash L)} \int_{\Lambda \backslash L} f d\mu,$$

where M is the centralizer of A in K , and μ is an L -invariant measure on $\Lambda \backslash L$.

A main ingredient of the proof of Theorem 2.5 is the work of N. Shah [S] (see Theorem 6.2 below) on the distribution in $\Lambda \backslash L$ of translates yUg as $g \rightarrow \infty$ for a subset $U \subset K$. Shah's result is based on Ratner's classification of measures invariant under unipotent flows [R1] and the work of Dani and Margulis on behavior of unipotent flows [DM]. Implementation of Shah's theorem in our setting is based on the fundamental property of the Furstenberg boundary \mathcal{B} of G : every regular element in a positive Weyl chamber acts on an open subset of full measure in \mathcal{B} as a contraction.

Remark 2.6 (on the rate of convergence). The method of the proof of Theorem 6.2 in [S] does not give any estimate on the rate of convergence. In the case when $L = G$ and $U = K$, Theorem 6.2 was proved by Eskin and McMullen in [EM]. The latter proof is based on the decay of the matrix coefficients of the quasi-regular representation of G on $L^2(\Gamma \backslash G)$ and provides an estimate on the rate of convergence. Combining the strong wavefront lemma (Theorem 2.1) with the method from [EM], we can derive an estimate for the rate of convergence in Theorem 2.5 when $L = G$ provided that one knows the rate of decay of matrix coefficients of $L^2(\Gamma \backslash G)$. In this case, it is also possible to obtain rates of convergence for the theorems stated in the introduction. We hope to address this problem in a sequel paper.

2.3. Equidistribution of lattice points in bisectors. For $\Omega_1, \Omega_2 \subset K$, and $g \in G$, we define

$$G_T(g, \Omega_1, \Omega_2) = \{h \in G : h \in g^{-1}\Omega_1 A^+ \Omega_2, d(K, Kgh) < T\}.$$

Using the strong wavefront lemma (Theorem 2.1 (2)), Theorem 1.6 is reduced to showing that the sets $G_T(g, \Omega_1, \Omega_2)$ are equidistributed in $\Gamma \backslash G$ in the sense of Theorem 2.5 for any Borel subsets $\Omega_1, \Omega_2 \subset K$ with boundaries of measure zero.

3. CARTAN DECOMPOSITION AND THE STRONG WAVEFRONT LEMMA

Let G be a connected noncompact semisimple Lie group with finite center, K a maximal compact subgroup of G , and $G = K \exp(\mathfrak{p})$ the Cartan decomposition determined by K . A split Cartan subgroup A with respect to K is a maximal connected abelian subgroup of G contained in $\exp(\mathfrak{p})$. It is well-known that two split Cartan subgroups with respect to K are conjugate to each other by an element of K . Fix a split Cartan subgroup A of G (with respect to K) with the set of positive roots Φ^+ and the positive Weyl chamber

$$A^+ = \{a \in A : \alpha(\log a) \geq 0 \text{ for all } \alpha \in \Phi^+\}.$$

Set $\mathfrak{a} = \log(A)$ and $\mathfrak{a}^+ = \log(A^+)$. Let M be the centralizer of A in K . Note that M is finite if and only if G is real split.

The following lemma is well-known (see, for example, [K, Ch. V]).

Lemma 3.1. (Cartan decomposition) *For every $g \in G$, there exist a unique element $\mu(g) \in \log(A^+)$ such that $g \in K \exp(\mu(g))K$. Moreover, if $k_1 a k_2 = k'_1 a k'_2$ for some a in the interior of A^+ , then there exists $m \in M$ such that $k_1 = k'_1 m$, $k_2 = m^{-1} k'_2$.*

Denote by d an invariant Riemannian metric on the symmetric space $K \backslash G$.

Lemma 3.2. *For every a_1 and a_2 in the interior of A^+ and $k \in K$,*

$$d(Ka_1, Ka_2) \leq d(Ka_1 k, Ka_2).$$

Proof. Let $a_i = \exp(H_i)$ for $H_i \in \mathfrak{a}^+$, $i = 1, 2$. Then $Ka_1k = K \exp(\text{Ad}(k^{-1})H_1)$. Applying the cosine inequality (see [H, Corollary I.13.2]) to the geodesic triangle with vertices Ke , Ka_1k , and Ka_2 , we obtain

$$\begin{aligned} d(Ka_1k, Ka_2)^2 &\geq d(K, Ka_1k)^2 + d(K, Ka_2)^2 - 2d(K, Ka_1k)d(K, Ka_2) \cos \alpha \\ &= \|H_1\|^2 + \|H_2\|^2 - 2\|H_1\|\|H_2\| \cos \alpha, \end{aligned}$$

where α is the angle at the vertex Ke . Since

$$\cos \alpha = \frac{\langle \text{Ad}(k^{-1})H_1, H_2 \rangle}{\|H_1\| \cdot \|H_2\|}$$

and by Lemma 3.3 below,

$$\langle \text{Ad}(k^{-1})H_1, H_2 \rangle \leq \langle H_1, H_2 \rangle,$$

it follows that

$$d(Ka_1k, Ka_2)^2 \geq \|H_1\|^2 + \|H_2\|^2 - 2\langle H_1, H_2 \rangle = \|H_1 - H_2\|^2 = d(Ka_1, Ka_2)^2.$$

The lemma is proved. \square

Lemma 3.3. *For any H_1 and H_2 in the interior of \mathfrak{a}^+ and for any $k \in K$,*

$$\langle H_1, H_2 \rangle \geq \langle \text{Ad}(k)H_1, H_2 \rangle.$$

Proof. By [H, Proposition VIII.5.2] and its proof, every G -invariant positive definite form on $K \backslash G$ is of the form $\sum_i \alpha_i B_i$, where B_i 's are the Killing forms of the simple factors of G and $\alpha_i > 0$. Thus, it is sufficient to consider the case when G is simple, and the Riemannian metric is given by the Killing form B of G .

Define the function $f(k) = \langle \text{Ad}(k)H_1, H_2 \rangle$ on K . Let $k_0 \in K$ be a point where f attains its maximum. For every $Z \in \text{Lie}(K)$,

$$\begin{aligned} 0 &= \left. \frac{d}{dt} \right|_{t=0} f(k_0 e^{tZ}) = \left. \frac{d}{dt} \right|_{t=0} B(\text{Ad}(k_0) \text{Ad}(e^{tZ})H_1, H_2) \\ &= B \left(\text{Ad}(k_0) \left. \frac{d}{dt} \right|_{t=0} (\text{Ad}(e^{tZ})H_1), H_2 \right) = B(\text{Ad}(k_0)(\text{ad}(Z)H_1), H_2) \\ &= B(\text{Ad}(k_0)[Z, H_1], H_2) = B(\text{Ad}(k_0)Z, [\text{Ad}(k_0)H_1, H_2]). \end{aligned}$$

This shows that $[\text{Ad}(k_0)H_1, H_2] \perp \text{Lie}(K)$. Since $[\text{Ad}(k_0)H_1, H_2] \in \text{Lie}(K)$ and the restriction of B to $\text{Lie}(K)$ is negative definite, it follows that $[\text{Ad}(k_0)H_1, H_2] = 0$. Therefore, $\text{Ad}(k_0)H_1 \in \mathfrak{a}$. Since the Weyl group W acts transitively on the set of Weyl chambers in A and K contains all representatives of the Weyl group, there exists an element $w \in K$ that normalizes A such that $\text{Ad}(w^{-1}k_0)H_1 \in \mathfrak{a}^+$. Since H_1 is in the interior of \mathfrak{a}^+ , it follows from the uniqueness of the Cartan decomposition (Lemma 3.1) that $\text{Ad}(k_0)H_1 = \text{Ad}(w)H_1$. It is easy to see [H, p. 288] that $\|\text{Ad}(w)H_1 - H_2\|$, $w \in W$, achieves its minimum at $w = e$. This implies that $\langle \text{Ad}(w)H_1, H_2 \rangle$ is maximal for $w = e$ and finishes the proof. \square

Proposition 3.4. *Let \mathcal{C} be a closed subset contained in A^+ with a positive distance from the walls of A^+ . Then for any neighborhood U_0 of e in K , there exists $\varepsilon > 0$ such that for any $a \in \mathcal{C}$,*

$$\{k \in K : d(Kak, Ka) < \varepsilon\} \subset MU_0.$$

Proof. Denote by Π the set of simple roots corresponding to the Weyl chamber A^+ . Without loss of generality, we may assume that

$$\mathcal{C} = \{a \in A^+ : \alpha(\log a) \geq C \text{ for all } \alpha \in \Pi\}$$

for some $C > 0$. Suppose that in contrary there exist sequences $\{a_i\} \subset \mathcal{C}$ and $\{k_i\} \subset K$ such that $d(Ka_i k_i, Ka_i) \rightarrow 0$ as $i \rightarrow \infty$, and no limit points of $\{k_i\}$ are contained in M . Passing to a subsequence, we may assume that $k_i \rightarrow k_0$ as $i \rightarrow \infty$ for some $k_0 \in K - M$, and for every $\alpha \in \Phi$, the sequence $\{\alpha(\log a_i)\}$ is either bounded or divergent. Set

$$\begin{aligned} \Phi_{\pm} &= \{\alpha \in \Phi : \alpha(\log a_i) \rightarrow \pm\infty \text{ as } i \rightarrow \infty\}, \\ \Phi_0 &= \{\alpha \in \Phi : \{\alpha(\log a_i)\} \text{ is bounded}\}. \end{aligned}$$

Let P^+ be the standard parabolic subgroup associated to $\Pi - \Phi_+$ and P^- the standard opposite parabolic subgroup for P^+ . Note that

$$P^- = \{g \in G : \{a_i g a_i^{-1}\} \text{ is bounded}\}.$$

Denote by U^+ and U^- the unipotent radicals of P^+ and P^- respectively. Set $Z = P^+ \cap P^-$, so that $P^{\pm} = ZU^{\pm}$. It is easy to see that $P^{\pm} \cap K \subset Z$. Denote by \mathfrak{g} , \mathfrak{u}^+ , \mathfrak{u}^- , \mathfrak{z} the corresponding Lie algebras and by \mathfrak{g}_{α} , $\alpha \in \Phi$, the root subspaces in \mathfrak{g} . We have

$$(3.5) \quad \mathfrak{u}^{\pm} = \bigoplus_{\alpha \in \Phi_{\pm}} \mathfrak{g}_{\alpha} \quad \text{and} \quad \mathfrak{z} = \bigoplus_{\alpha \in \Phi_0} \mathfrak{g}_{\alpha}.$$

Step 1: We claim that $k_0 \in Z$.

There is an embedding $\pi : G/C_G \rightarrow \mathrm{SL}_d(\mathbb{R})$, where C_G is the center of G , such that $\pi(AC_G)$ is contained in the group of diagonal matrices. We have

$$\pi(a_i k_i a_i^{-1})_{st} = \pi(a_i)_{ss} \pi(a_i)_{tt}^{-1} \cdot \pi(k_i)_{st}, \quad s, t = 1, \dots, d.$$

Passing to a subsequence, we may assume that for each (s, t) , the sequence $\{\pi(a_i)_{ss} \pi(a_i)_{tt}^{-1}\}$ is either bounded or divergent. If the sequence $\{\pi(a_i)_{ss} \pi(a_i)_{tt}^{-1}\}$ is divergent, then $\pi(k_i)_{st} \rightarrow 0$ as $i \rightarrow \infty$. Thus, $\pi(k_0)_{st} = 0$ for every pair (s, t) such that $\{\pi(a_i)_{ss} \pi(a_i)_{tt}^{-1}\}$ is divergent. It follows that $\pi(a_i k_0 a_i^{-1})$ is bounded. Since the center of G is finite, this proves that the sequence $\{a_i k_0 a_i^{-1}\}$ is bounded. Thus, $k_0 \in P^-$. Since $P^- \cap K \subset Z$, the claim follows.

Step 2: We claim that $d(Ka_i k_0 a_i^{-1}, K) \rightarrow 0$ as $i \rightarrow \infty$.

Write $k_i = k_0 l_i$ where $l_i \in K$ and $l_i \rightarrow e$ as $i \rightarrow \infty$. Then

$$(3.6) \quad d(Ka_i k_i, Ka_i) = d(Ka_i k_0 a_i^{-1}, Ka_i l_i^{-1} a_i^{-1}) \rightarrow 0 \text{ as } i \rightarrow \infty.$$

Thus, the sequence $\{a_i l_i^{-1} a_i^{-1}\}$ is bounded, and hence, we may assume that it converges. Since $l_i^{-1} \rightarrow e$ as $i \rightarrow \infty$ and $\mathfrak{g} = \mathfrak{u}^- \oplus \mathfrak{z} \oplus \mathfrak{u}^+$, we obtain that $l_i^{-1} = u_i^- z_i u_i^+$ for some $u_i^- \in U^-$, $z_i \in Z$, and $u_i^+ \in U^+$ such that $u_i^- \rightarrow e$, $z_i \rightarrow e$, and $u_i^+ \rightarrow e$ as $i \rightarrow \infty$. It follows from (3.5) that $a_i u_i^- a_i^{-1} \rightarrow e$ and $a_i z_i a_i^{-1} \rightarrow e$ as $i \rightarrow \infty$. Hence, $a_i l_i^{-1} a_i^{-1} \rightarrow u^+$ as $i \rightarrow \infty$ for some $u^+ \in U^+$. On the other hand, by Step 1, passing to a subsequence, we get $a_i k_0 a_i^{-1} \rightarrow z$ as $i \rightarrow \infty$ for some $z \in Z$. Thus, by (3.6), $z^{-1} u^+ \in K$ as $i \rightarrow \infty$. Since $P^+ \cap K \subset Z$, we deduce that $u^+ = e$. Hence, $a_i l_i^{-1} a_i \rightarrow e$ as $i \rightarrow \infty$, and the claim follows from (3.6).

Step 3: We claim that $a_i k_0 a_i^{-1} = d_i k_0 d_i^{-1}$ for some bounded sequence $\{d_i\} \subset \mathcal{C}$.

Recall that the system of simple roots Π is a basis of the dual space of the Lie algebra of A . Hence, we may write $a_i = b_i c_i$ where $b_i, c_i \in A^+$ such that $\alpha(\log b_i) = 0$ for every $\alpha \in \Pi \cap \Phi_+$ and $\alpha(\log c_i) = 0$ for every $\alpha \in \Pi - \Phi_+$. Then c_i commutes with Z , and since $\alpha(b_i) = \alpha(a_i)$ for every $\alpha \in \Pi - \Phi_+$, the sequence $\{b_i\}$ is bounded. Let $c_0 \in A^+$ be such that $\alpha(\log c_0) = C$ for every $\alpha \in \Pi \cap \Phi_+$ and $\alpha(\log c_0) = 0$ for every $\alpha \in \Pi - \Phi_+$. Then $d_i = b_i c_0 \in \mathcal{C}$ and $a_i k_0 a_i^{-1} = d_i k_0 d_i^{-1}$ as required.

Taking a subsequence, we obtain that $d_i \rightarrow d_0$ as $i \rightarrow \infty$ for some $d_0 \in \mathcal{C}$ and $d_0 k_0 d_0^{-1} \in K$. This implies that $d_0 k_0 = k d_0$ for some $k \in K$. Since d_0 lies in the interior of A^+ , it follows from the uniqueness properties of the Cartan decomposition (Lemma 3.1) that $k_0 = k \in M$. This is a contradiction. \square

Theorem 3.7 (The strong wavefront lemma). *Let \mathcal{C} be a closed subset contained in A^+ with a positive distance from the walls of A^+ . Then for any neighborhoods U of e in K and V of e in A , there exists a neighborhood \mathcal{O} of e in G such that*

$$\mathcal{O}g \cup g\mathcal{O} \subset (k_1 U)(aVM)(k_2 U)$$

for any $g = k_1 a k_2 \in KCK$.

Proof. Without loss of generality, we may assume that $\mathcal{C} = A^+(C)$ for some $C > 0$, where

$$A^+(C) = \{a \in A^+ : \alpha(\log a) \geq C \text{ for all } \alpha \in \Phi^+\}.$$

Replacing V by a smaller subset if necessary, we may assume that $A^+(C)V$ is contained in the interior of $A^+(C/2)$.

Step 1: We claim that there exists an ε -neighborhood \mathcal{O} of e in G such that

$$\mathcal{O}g \subset KaVK \quad \text{for all } g = k_1 a k_2 \in KA^+(C)K.$$

We take $\varepsilon > 0$ to be sufficiently small such that the ε -neighborhood of e in A is contained in V . Let \mathcal{O} be the ε -neighborhood of e in G . For $h \in \mathcal{O}a$,

write $h = k'_1 b k'_2 \in KA^+K$. Then $d(Kbk'_2 a^{-1}, K) = d(Kbk'_2, Ka) < \varepsilon$. Since $d(Kb, Ka) \leq d(Kbk'_2, Ka)$ by Lemma 3.2, we have

$$(3.8) \quad d(Kb, Ka) < \varepsilon$$

and hence, $h = k'_1 b k'_2 \in KVaK$. This shows that $\mathcal{O}a \subset KaVK$. Since \mathcal{O} is K -invariant,

$$\mathcal{O}k_1 a k_2 \subset \mathcal{O}a k_2 \subset KaVK$$

This proves the claim.

Step 2: We claim that there exists an ε -neighborhood \mathcal{O} of e in G such that

$$(3.9) \quad \mathcal{O}g \subset KA^+(k_2U) \quad \text{for all } g = k_1 a k_2 \in KA^+(C)K.$$

Let U_0 be a neighborhood of e in K such that $kU_0k^{-1} \subset U$ for all $k \in K$. Choose $\varepsilon > 0$ that satisfies Step 1 above, and Proposition 3.4 holds with respect to 2ε , U_0 , and $A^+(C/2)$. Take \mathcal{O} to be the ε -neighborhood of e in G . Then for any $h = k'_1 b k'_2 \in \mathcal{O}a$, we have by (3.8),

$$d(Kbk'_2, Kb) \leq d(Kbk'_2, Ka) + d(Kb, Ka) < 2\varepsilon.$$

Since $\mathcal{O}a \subset KA^+(C/2)K$, by Proposition 3.4, this implies that $k'_2 \in MU_0$, and hence $\mathcal{O}a \subset KA^+MU_0$. Thus,

$$\mathcal{O}(k_1 a k_2) \subset KA^+MU_0 k_2 \subset KA^+M(k_2U).$$

Step 3: We claim that there exists a neighborhood \mathcal{O} of e in G such that

$$\mathcal{O}g \subset (k_1U)A^+K \quad \text{for all } g = k_1 a k_2 \in KA^+(C)K.$$

Recall that by the Iwasawa decomposition, $G = KAN$, where N is the subgroup of G corresponding to the sum of the positive root spaces. Since the Weyl group acts transitively on the sets of simple roots, there exists an element $w \in K$ that normalizes A such that $wA^+(C)w^{-1} = (A^+(C))^{-1}$.

Let U_0 be a neighborhood of e in K such that $U_0(wU_0^{-1}w^{-1}) \subset U$. Let \mathcal{O} be the neighborhood from Step 2 with respect to U_0 and \mathcal{O}_1 a neighborhood of e in AN such that $c^{-1}\mathcal{O}_1c \subset \mathcal{O}$ for all $c \in A^+$. Since $w \in K$, $w\mathcal{O}w^{-1} = \mathcal{O}$. Conjugating (3.9) (with respect to U_0) by w , we have for any $b \in wA^+(C)w^{-1}$,

$$\mathcal{O}b \subset K(A^+)^{-1}(wU_0w^{-1}).$$

For $a \in A^+(C)$, $a^{-1} \in wA^+(C)w^{-1}$ and hence

$$(3.10) \quad a^{-1}\mathcal{O}_1 \subset \mathcal{O}a^{-1} \subset K(A^+)^{-1}(wU_0w^{-1}).$$

By taking the inverse of (3.10),

$$(3.11) \quad \mathcal{O}_1^{-1}a \subset (wU_0^{-1}w^{-1})A^+K.$$

Since the product map $K \times AN \rightarrow G$ is a diffeomorphism, $U_0\mathcal{O}_1^{-1}$ is a neighborhood of e in G . Therefore, there exists a neighborhood \mathcal{O}_2 of e in G such that $k^{-1}\mathcal{O}_2k \subset U_0\mathcal{O}_1^{-1}$ for all $k \in K$. Then by (3.11),

$$\mathcal{O}_2k_1ak_2 \subset k_1U_0\mathcal{O}_1^{-1}ak_2 \subset k_1U_0(wU_0^{-1}w^{-1})A^+K \subset k_1UA^+K.$$

This finishes the proof of Step 3.

By the above three steps, we obtain a neighborhood \mathcal{O}_1 of e in G such that for all $g = k_1ak_2 \in KA^+(C)K$,

$$\mathcal{O}_1g \subset KaVK \cap KA^+(k_2U) \cap (k_1U)A^+K.$$

By the uniqueness the Cartan decomposition (Lemma 3.1),

$$KaVK \cap KA^+(k_2U) \cap (k_1U)A^+K = (k_1U)(aVM)(k_2U).$$

Hence, we have shown that for every neighborhood U of e in K and every neighborhood V of e in A , there exists a neighborhood \mathcal{O}_1 of e in G such that

$$(3.12) \quad \mathcal{O}_1g \subset (k_1U)(aVM)(k_2U) \quad \text{for all } g = k_1ak_2 \in KA^+(C)K.$$

Let U_0 be a neighborhood of e in K such that $kU_0^{-1}k^{-1} \subset U$ for every $k \in K$ and V_0 a neighborhood of e in A such that $wV_0^{-1}w^{-1} \subset V$, where w is the element of K defined in Step 3. Let \mathcal{O}_2 be a neighborhood of e in G such that

$$\mathcal{O}_2g \subset (k_1U_0)(aV_0M)(k_2U_0) \quad \text{for all } g = k_1ak_2 \in KA^+(C)K.$$

By taking the inverse, we have

$$g\mathcal{O}_2^{-1} \subset (U_0^{-1}k_1)(aV_0^{-1}M)(U_0^{-1}k_2) \quad \text{for all } g = k_1ak_2 \in KA^+(C)^{-1}K.$$

Conjugating by w , we get

$$g(w\mathcal{O}_2^{-1}w^{-1}) \subset (k_1U)(aVM)(k_2U) \quad \text{for all } g = k_1ak_2 \in KA^+(C)K.$$

Set $\mathcal{O} = \mathcal{O}_1 \cap (w\mathcal{O}_2^{-1}w^{-1})$ to finish the proof. \square

4. CONTRACTIONS ON G/B

Let G be a connected noncompact semisimple Lie group with finite center, K a maximal compact subgroup of G , and B a maximal connected split solvable subgroup of G . Let N be the unipotent radical of B . The normalizer P of N in G is the unique minimal parabolic subgroup of G containing B .

Lemma 4.1. *There exists a split Cartan subgroup A of G with respect to K and an ordering on the root system Φ such that*

- $G = KA^+K$.
- $B = AN$.
- N is the subgroup generated by all positive root subgroups of G with respect to A .
- $P = MAN$, where M is the centralizer of A in K and $M = K \cap P$.

Proof. Take any split Cartan subgroup A_0 and a Weyl chamber A_0^+ such that the Cartan decomposition $G = KA_0^+K$ holds. Set $P_0 = M_0A_0N_0$, where M_0 is the centralizer of A_0 in K , and N_0 is the subgroup generated by all positive root subgroups of G with respect to A_0 . Note that P_0 is a minimal parabolic subgroup of G (see [W, Sec. 1.2.3]). By the Iwasawa decomposition, $G = KP_0$. Since all minimal parabolic subgroups are conjugate to each other ([loc.cit.]), there exists $k \in K$ such that $P = kP_0k^{-1}$. Set $A = kA_0k^{-1}$, $A^+ = kA_0^+k^{-1}$, and $M = kM_0k^{-1}$. It is clear that $N = kN_0k^{-1}$. Since AN is normal in P , it is the unique maximal connected split solvable subgroup in P . Thus, $B = AN$. \square

Let A be a split Cartan subgroup as in Lemma 4.1. Denote by M be the centralizer of A in K and by N^- the subgroup generated by all negative root subgroups of G . One can check that the map

$$N^- \times M \rightarrow G/B : (n, m) \mapsto nmB$$

is a diffeomorphism onto its image. Moreover, it follows from the properties of Bruhat decomposition that $G/B - \pi(N^- \times M)$ is a finite union of closed submanifolds of smaller dimensions. On the other hand, by the Iwasawa decomposition, the map

$$K \rightarrow G/B : k \mapsto kB.$$

is a diffeomorphism. Thus, we have a map $N^- \times M \rightarrow K$. Since M normalizes B , this map is right M -equivariant. Denote by $S \subset K$ the submanifold which is the image of $N^- \times \{e\}$ under this map.

Let ν be the probability Haar measure on K and τ the probability Haar measure on M . Then $\nu = \sigma \otimes \tau$ for some finite smooth measure σ on S (see [W, p. 73]), i.e.,

$$(4.2) \quad \int_K f d\nu = \int_M \int_S f(sm) d\sigma(s) d\tau(m), \quad f \in C(K).$$

For $C > 0$, put

$$A^+(C) = \{a \in A : \alpha(\log a) \geq C \text{ for all } \alpha \in \Phi^+\}.$$

Lemma 4.3. *Let Ω be a Borel subset of K , $s \in S$, $m \in M$, and $k \in K$. For $a \in A$, $r \in K$, and $U \subset K$, define*

$$\Omega_r(k, a, U) = \{l \in \Omega : lkB \in a^{-1}r^{-1}UB\}.$$

- (1) *Let $VW \subset K$ be an open neighborhood of s where $V \subset S$ and $W \subset M$ are open subsets. Then for every $\varepsilon > 0$, there exists $C > 0$ such that*

$$\nu((\Omega \cap m^{-1}SWk^{-1}) - \Omega_{sm}(k, a, VW)) < \varepsilon \text{ for all } a \in A^+(C).$$

- (2) *Let $U \subset K$ be a Borel subset such that $s \notin \overline{UM}$. Then for every $\varepsilon > 0$, there exists $C > 0$ such that*

$$\nu(\Omega_{sm}(k, a, U)) < \varepsilon \text{ for all } a \in A^+(C).$$

Proof. Note that $\Omega_{sm}(k, a, U) = m^{-1} \cdot (m\Omega)_s(k, a, U)$. Hence, replacing Ω by $m\Omega$, we may assume that $m = e$. Also, $\Omega_s(k, a, U) = (\Omega k)_s(e, a, U) \cdot k^{-1}$. Thus, we may assume that $k = e$.

The proof is based on the observation that elements in the interior of A^+ act on N^-B as contractions (cf. [Z, Proposition 8.2.5]). Denote by π the map $\pi : N^- \rightarrow G/B : n \mapsto nB$. Note that for $a \in A$ and $V_0 \subset S$,

$$(4.4) \quad \pi^{-1}(a^{-1}V_0B) = a^{-1}\pi^{-1}(V_0B)a.$$

Let D be a compact set in S such that $\sigma(S - D) < \varepsilon/2$.

Suppose that $s \in VW$. Let $W_0 \subset W$ be a compact set such that $\tau(W - W_0) < \varepsilon/2$. Then $s^{-1}VW$ is a neighborhood of W_0 . By uniform continuity, there exists a neighborhood V_0 of e in S such that $V_0W_0 \subset s^{-1}VW$. For each $a \in A^+$, the map $N^- \rightarrow N^- : x \rightarrow a^{-1}xa$ expands any neighborhood of e at least by the factor of $\min_{\alpha \in \Phi^+} e^{\alpha(\log a)}$. Hence, there exists $C > 0$ such that $\pi^{-1}(DB) \subset a^{-1}\pi^{-1}(V_0B)a$ for all $a \in A^+(C)$. Then by (4.4), $DB \subset a^{-1}V_0B$. It follows that $DW_0B \subset a^{-1}V_0W_0B$ and

$$\begin{aligned} \nu(\Omega_s(e, a, VW)) &\geq \nu(\Omega \cap DW_0) \geq \nu(\Omega \cap SW) - \nu((S - D)M) - \nu(S(W - W_0)) \\ &\geq \nu(\Omega \cap SW) - \varepsilon. \end{aligned}$$

This proves the first part of the lemma.

To prove the second part, we observe that there exists an open subset $V \subset S$ such that $s \in V \subset K - \overline{UM}$. Since $\nu(m^{-1}SMk^{-1}) = 1$, it suffices to apply the first part with $W = M$. \square

Lemma 4.5. *Let $W \subset M$ and $\Omega \subset K$ be Borel subsets. Then every $k \in K$,*

$$\int_M \nu(\Omega \cap m^{-1}SWk^{-1}) d\tau(m) = \nu(\Omega)\tau(W).$$

Proof. Without loss of generality, $k = e$. Since M normalizes N^- , we have $m^{-1}Sm \subset S$. Since $\nu(K - SM) = 0$, we may assume that $\Omega \subset SM$. Moreover, it suffices to prove the lemma for $\Omega = \Omega_S\Omega_M$ with Borel sets $\Omega_S \subset S$ and $\Omega_M \subset M$. By (4.2),

$$\nu(\Omega \cap m^{-1}SW) = \sigma(\Omega_S)\tau(\Omega_M \cap m^{-1}W).$$

For $Y \subset M$, denote by χ_Y the characteristic function of Y . We have

$$\begin{aligned} \int_M \tau(\Omega_M \cap m^{-1}W) d\tau(m) &= \int_M \int_M \chi_{\Omega_M}(l)\chi_{m^{-1}W}(l) d\tau(l)d\tau(m) \\ &= \int_M \chi_{\Omega_M}(l) \left(\int_M \chi_{Wl^{-1}}(m) d\tau(m) \right) d\tau(l) = \tau(\Omega_M)\tau(W). \end{aligned}$$

Since $\nu(\Omega) = \sigma(\Omega_S)\tau(\Omega_M)$, this proves the lemma. \square

5. VOLUME ESTIMATES

Let G be a connected noncompact semisimple Lie group with finite center and $G = KA^+K$ a Cartan decomposition. Let dk denote the probability Haar measure on K , dt the Lebesgue measure on the Lie algebra \mathfrak{a} of A , and da the Haar measure on A derived from dt via the exponential map. We denote by m the Haar measure on G which is normalized so that for any $f \in C_c(G)$,

$$(5.1) \quad \int_G f dm = \int_K \int_{A^+} \int_K f(k_1 a k_2) \xi(\log a) dk_1 da dk_2,$$

where

$$(5.2) \quad \xi(t) = \prod_{\alpha \in \Phi^+} \sinh(\alpha(t))^{m_\alpha}, \quad t \in \mathfrak{a},$$

and m_α is the dimension of the root subspace corresponding to α . In particular, for any measurable subset $D \subset A^+$, we have

$$(5.3) \quad \text{Vol}(KDK) = \int_D \xi(\log a) da.$$

Let $\|\cdot\|$ be a Euclidean norm on \mathfrak{a} , that is, for some basis v_1, \dots, v_n of \mathfrak{a} , $\|\sum_i c_i v_i\| = \sqrt{\sum_i c_i^2}$. We assume that $\|\cdot\|$ is invariant under the Weyl group action. For instance, $\|\cdot\|$ can be taken to be the norm induced from an invariant Riemannian metric d on the symmetric space $K \backslash G$, i.e.,

$$\|t\| = d(K, K \exp(t)) \quad \text{for } t \in \mathfrak{a}.$$

For $T > 0$ and $\mathfrak{t} \subset \mathfrak{a}$, set

$$\mathfrak{t}_T = \{t \in \mathfrak{t} : \|t\| < T\}.$$

Let $\rho = \frac{1}{2} \sum_{\alpha \in \Phi^+} m_\alpha \alpha$. One can check that the maximum of ρ on $\bar{\mathfrak{a}}_1$ is achieved at a unique point contained in the interior of \mathfrak{a}^+ , which we call the *barycenter* of \mathfrak{a}^+ .

We present a simple derivation of the asymptotics of the volume of Riemannian balls in a symmetric space of noncompact type (see also [Kn, Theorem 6.2]).

Lemma 5.4. *Let \mathfrak{q} be a convex cone in \mathfrak{a}^+ centered at the origin that contains the barycenter of \mathfrak{a}^+ in its interior. Then for some $C > 0$ (independent of \mathfrak{q}),*

$$\int_{\mathfrak{q}_T} \xi(t) dt \sim_{T \rightarrow \infty} C \cdot T^{(r-1)/2} e^{\delta T},$$

where $r = \mathbb{R}\text{-rank}(G) = \dim A$ and $\delta = \max\{2\rho(t) : t \in \bar{\mathfrak{a}}_1\}$. In particular,

$$\int_{\mathfrak{q}_T} \xi(t) dt \sim_{T \rightarrow \infty} \int_{\mathfrak{a}_T^+} \xi(t) dt.$$

Proof. We have

$$\int_{\mathfrak{q}_T} \xi(t) dt = \frac{1}{2^{|\Phi^+|}} \int_{\mathfrak{q}_T} e^{2\rho(t)} dt + \text{“other terms”},$$

where the “other terms” are linear combinations of integrals of the form $\int_{\mathfrak{q}_T} e^{\lambda(t)} dt$ such that $2\rho - \lambda = \sum_{\alpha \in \Phi^+} n_\alpha \alpha$ for some $n_\alpha \geq 0$. In particular, $2\rho > \lambda$ in the interior of the Weyl chamber \mathfrak{a}^+ . Since the maximum of 2ρ in $\bar{\mathfrak{a}}_1$ is achieved in the interior of \mathfrak{q} ,

$$\max\{2\rho(t) : t \in \bar{\mathfrak{q}}_1\} > \max\{\lambda(t) : t \in \bar{\mathfrak{q}}_1\} \stackrel{def}{=} \delta'.$$

Thus,

$$\int_{\mathfrak{q}_T} e^{\lambda(t)} dt \leq e^{\delta'T} \text{Vol}(\mathfrak{q}_T) \ll e^{\delta'T} T^r = o(e^{\delta'T})$$

as $T \rightarrow \infty$. It remains to show that for some $C > 0$ independent of \mathfrak{q} ,

$$(5.5) \quad \int_{\mathfrak{q}_T} e^{2\rho(t)} dt \sim_{T \rightarrow \infty} C \cdot T^{(r-1)/2} e^{\delta'T}.$$

Making a change variables and decomposing \mathfrak{q}_1 into slices parallel to the hyperplane $\{2\rho = 0\}$, we have

$$\int_{\mathfrak{q}_T} e^{2\rho(t)} dt = T^r \int_{\mathfrak{q}_1} e^{2T\rho(t)} dt = T^r \int_0^\delta e^{Tx} \phi(x) dx,$$

where $\phi(x) = \text{Vol}_{r-1}(\mathfrak{q}_1 \cap \{2\rho = x\})$.

First, we show that for some $c_1 > 0$ independent of \mathfrak{q} ,

$$(5.6) \quad \phi(x) \sim_{x \rightarrow \delta^-} c_1 \cdot (\delta - x)^{(r-1)/2}.$$

We identify \mathfrak{a} with the set $\{(t_1, \dots, t_r) \in \mathbb{R}^r\}$ and denote by Q the positive quadratic form on \mathfrak{a} defined by the norm. After a linear change of variables, we may assume that $2\rho(t) = t_r$ and

$$Q(t_1, \dots, t_r) = \sum_{i=1}^{r-1} \alpha_i (t_i - \beta_i t_r)^2 + \alpha_r t_r^2$$

for some $\alpha_i > 0$ and $\beta_i \in \mathbb{R}$. It is clear that the maximum of $2\rho(t)$ on the set $\bar{\mathfrak{a}}_1 = \{t : Q(t) \leq 1\}$ is achieved when $t_i = \beta_i t_r$ for $i = 1, \dots, r-1$ and $\alpha_r t_r^2 = 1$. This implies that $\delta = \alpha_r^{-1/2}$. Since for x close to δ , the set $\mathfrak{q}_1^+ \cap \{2\rho = x\}$ is defined by the condition $Q(t_1, \dots, t_{r-1}, x) \leq 1$, we have

$$\begin{aligned} \phi(x) &= \text{Vol}_{r-1} \left(\left\{ (t_1, \dots, t_{r-1}) : \sum_{i=1}^{r-1} \alpha_i (t_i - \beta_i x)^2 \leq 1 - \alpha_r x^2 \right\} \right) \\ &= c_1 \cdot (1 - \alpha_r x^2)^{(r-1)/2} = c_1 \cdot (1 - x^2/\delta^2)^{(r-1)/2} \end{aligned}$$

for a constant $c_1 > 0$. This proves (5.6). Now by (5.6) and the L'Hopital's rule,

$$\beta(x) \stackrel{\text{def}}{=} \int_0^x \phi(\delta - u) du \sim_{x \rightarrow 0^+} c_2 \cdot x^{(r+1)/2}$$

for some $c_2 > 0$. Thus, by the Abelian theorem (see [Wi, Corollary 1a, p. 182]),

$$\begin{aligned} \int_0^\delta e^{Tx} \phi(x) dx &= e^{\delta T} \int_0^\delta e^{-Tx} \phi(\delta - x) dx = e^{\delta T} \int_0^\infty e^{-Tx} d\beta(x) \\ &\sim_{T \rightarrow \infty} c_3 \cdot T^{-(r+1)/2} e^{\delta T} \end{aligned}$$

for some $c_3 > 0$. It is clear that c_3 is independent of \mathfrak{q} . This proves (5.5) and the proposition. \square

For $T > 0$ and $R \subset G$, define

$$(5.7) \quad R_T = \{r \in R : d(K, Kr) < T\}.$$

Since $G_T = KA_T^+K$, combining the previous lemma and (5.3), we deduce

Corollary 5.8. *For some $C > 0$,*

$$\text{Vol}(G_T) \sim_{T \rightarrow \infty} C \cdot T^{(r-1)/2} e^{\delta T}.$$

In particular, we have

Lemma 5.9. *For some functions $a(\varepsilon)$ and $b(\varepsilon)$ such that $a(\varepsilon) \rightarrow 1$ and $b(\varepsilon) \rightarrow 1$ as $\varepsilon \rightarrow 0^+$, we have*

$$a(\varepsilon) \leq \liminf_{T \rightarrow \infty} \frac{\text{Vol}(G_{T-\varepsilon})}{\text{Vol}(G_T)} \leq \limsup_{T \rightarrow \infty} \frac{\text{Vol}(G_{T+\varepsilon})}{\text{Vol}(G_T)} \leq b(\varepsilon).$$

For $T, C > 0$ and $R \subset A$, define

$$(5.10) \quad \begin{aligned} R(C) &= \{r \in R : \alpha(\log r) \geq C \text{ for all } \alpha \in \Phi^+\}, \\ R_T(C) &= R_T \cap R(C). \end{aligned}$$

Lemma 5.11. *Let $Q \subset A^+$ be a convex cone centered at the origin that contains the barycenter in its interior. Then for any fixed $C > 0$,*

$$\int_{Q_T(C)} \xi(\log a) da \sim_{T \rightarrow \infty} \int_{A_T^+} \xi(\log a) da.$$

In particular,

$$\text{Vol}(KA_T^+K) \sim_{T \rightarrow \infty} \text{Vol}(KA_T^+(C)K).$$

Proof. It suffices to prove the lemma when Q is contained in the interior of A^+ . Then there exists $T_0 > 0$ such that $Q_T(C) \supset Q_T - Q_{T_0}$ for all sufficiently large $T > 0$. Thus, the lemma follows from Lemma 5.4. \square

6. EQUIDISTRIBUTION OF SOLVABLE FLOWS

Let G be a connected noncompact semisimple Lie group with finite center which is realized as a closed subgroup of a Lie group L , Λ a lattice in L , and Q a closed subgroup of G that contains a maximal connected split solvable subgroup of G . In this section we investigate the distribution of orbits of Q in the homogeneous space $\Lambda \backslash L$.

Let K_0 be a maximal compact subgroup of G and A_0 a split Cartan subgroup of G with respect to K_0 . Then $G = K_0 A_0^+ K_0$ for any positive Weyl chamber A_0^+ in A_0 . Denote by d an invariant Riemannian metric on $K_0 \backslash G$. For $g \in G$, $R \subset G$, $\Omega \subset K_0$, and $T > 0$, define

$$R_T(g, \Omega) = \{r \in R : d(K_0, K_0gr) < T, r \in g^{-1}K_0A_0^+\Omega\}$$

and $R_T(g) = R_T(g, K_0)$. Note that

$$R_T(k_0g, \Omega) = R_T(g, \Omega) = g^{-1}(gR)_T(e, \Omega)$$

for any $k_0 \in K_0$.

The main result in this section is Theorem 6.1 on equidistribution of the sets $Q_T(g, \Omega)$ as $T \rightarrow \infty$. Let μ be the Haar measure on L such that $\mu(\Lambda \backslash L) = 1$, ν_0 the probability Haar measure on K_0 , and ρ a right invariant Haar measure on Q . Denote by G_n the product of all noncompact simple factors of G .

Theorem 6.1. *Suppose that for $y \in \Lambda \backslash L$, the orbit yG_n is dense in $\Lambda \backslash L$. Then for $g \in G$, any Borel subset $\Omega \subset K_0$ with boundary of measure zero and $f \in C_c(\Lambda \backslash L)$,*

$$\lim_{T \rightarrow \infty} \frac{1}{\rho(Q_T(g))} \int_{Q_T(g, \Omega)} f(yq^{-1}) d\rho(q) = \nu_0(M_0\Omega) \int_{\Lambda \backslash L} f d\mu.$$

The rest of this section is devoted to a proof of Theorem 6.1. We start by stating a theorem of Shah ([S, Corollary 1.2]). Recall that a sequence $\{g_i\} \subset G$ is called *strongly divergent* if for every projection π from G to its noncompact factor, $\pi(g_i) \rightarrow \infty$ as $i \rightarrow \infty$.

Theorem 6.2 (Shah). *Suppose that for $y \in \Lambda \backslash L$, the orbit yG_n is dense in $\Lambda \backslash L$. Let $\{g_i\} \subset G$ be a strongly divergent sequence. Then for any $f \in C_c(\Lambda \backslash L)$ and any Borel subset U in K_0 with boundary of measure zero,*

$$\lim_{i \rightarrow \infty} \int_U f(ykg_i) d\nu_0(k) = \nu_0(U) \int_{\Lambda \backslash L} f d\mu.$$

Remark 6.3. Although this theorem was stated in [S] only for the case $U = K_0$, its proof works equally well when U is an open subset of K_0 with boundary of measure zero, and approximating Borel sets by open sets, one can check that Theorem 6.2 holds in the above generality.

Let $K = g^{-1}K_0g$ and B be a maximal connected split solvable subgroup of G . By Lemma 4.1, there exists a split Cartan subgroup A with an ordering on the root system Φ such that $G = KA^+K$ and $B = AN$, where N is the subgroup generated by all positive root subgroups of A in G . Let M be the centralizer of A in K .

Let m be the Haar measure on G such that (5.1) holds with respect to K_0 and A_0^+ . It follows from the uniqueness of the Haar measure that for every $g \in G$, there exists $c_g > 0$ such that

$$(6.4) \quad \int_G f dm = c_g \int_{K_0} \int_B f(g^{-1}kgb) d\rho(b) d\nu_0(k), \quad f \in C_c(G).$$

In particular,

$$(6.5) \quad c_g \rho(B_T(g)) = \text{Vol}(G_T(g)) = \text{Vol}(G_T(e)).$$

We normalize ρ so that $c_e = 1$. Let ν be the probability Haar measure on K .

We use notations from Section 4. In particular, (4.2) holds.

Proposition 6.6. *Suppose that for $y \in \Lambda \backslash L$, the orbit yG_n is dense in $\Lambda \backslash L$. Let $U = VW$ be an open neighborhood of e in K where V is an open neighborhood of e in S , and W is an open neighborhood of e in M such that $\sigma(\partial V) = \tau(\partial W) = 0$ and Ω a Borel subset of K_0 such that $\nu_0(\partial\Omega) = 0$. Then for any $f \in C_c(\Lambda \backslash L)$,*

$$\frac{1}{\rho(B_T(g))} \int_U \int_{B_T(g, \Omega)} f(yb^{-1}k^{-1}) d\rho(b) dk \rightarrow \nu(U) \nu_0(M_0\Omega) \int_{\Lambda \backslash L} f d\mu$$

as $T \rightarrow \infty$.

Proof. Since both A_0 and gAg^{-1} are split Cartan subgroups with respect to K_0 , there exists $k_0 \in K_0$ such that $A_0 = k_0gAg^{-1}k_0^{-1}$. Hence, replacing g by k_0g , we may assume that $A = g^{-1}A_0g$.

Let d be the Riemannian metric on $K \backslash G$ induced from the metric on $K_0 \backslash G$ by the map $Kx \mapsto Kgx$, and notations A_T^+ , $A_T^+(C)$, and $A^+(C)$ be defined as in Section 5. For instance,

$$A_T^+ = \{a \in A^+ : d(K_0g, K_0ga) < T\}.$$

Since $KA_T^+K = g^{-1}G_T(e)g$, it follows from (5.3) and (6.5) that for every $T > 0$,

$$(6.7) \quad \rho(B_T(e)) = \int_{A_T^+} \xi(\log a) da$$

and by Lemma 5.11,

$$(6.8) \quad \int_{A_T^+ - A_T^+(C)} \xi(\log a) da = o(\rho(B_T(e))) \quad \text{as } T \rightarrow \infty.$$

Since $B_T(g, \Omega) = B_T(g, M_0\Omega)$, we may assume without loss of generality that $M_0\Omega = \Omega$.

We have

$$\begin{aligned} UB_T(g, \Omega) &= g^{-1}K_0A_{0T}^+(e)\Omega \cap UB = KA_T^+(g^{-1}\Omega g)g^{-1} \cap UB \\ &= \{k_1ak_2g^{-1} : k_1 \in K, a \in A_T^+, k_2 \in \Omega_{k_1}(a)\}, \end{aligned}$$

where

$$\Omega_{k_1}(a) = \{k_2 \in g^{-1}\Omega g : k_2g^{-1}B \in a^{-1}k_1^{-1}UB\}.$$

By (5.1) and (6.4),

$$\begin{aligned} (6.9) \quad c_g \int_U \int_{B_T(g, \Omega)} f(yb^{-1}k^{-1}) d\rho(b) d\nu(k) &= \int_{UB_T(g, \Omega)} f(yx^{-1}) dm(x) \\ &= \int_{k_1ak_2g^{-1} \in UB_T(g, \Omega)} f(ygk_2^{-1}a^{-1}k_1^{-1}) \xi(\log a) d\nu(k_2) dad\nu(k_1) \\ &= \int_{k_1 \in K} \int_{a \in A_T^+} \int_{k_2 \in \Omega_{k_1}(a)} f(ygk_2^{-1}a^{-1}k_1^{-1}) \xi(\log a) d\nu(k_2) dad\nu(k_1). \end{aligned}$$

Step 1: We claim that for every $m \in M$,

$$\begin{aligned} (6.10) \quad \lim_{T \rightarrow \infty} \frac{1}{\rho(B_T(e))} \int_V \int_{A_T^+} \int_{\Omega_{sm}(a)} f(ygk^{-1}a^{-1}(sm)^{-1}) \xi(\log a) d\nu(k) dad\sigma(s) \\ = \sigma(V) \cdot \nu(g^{-1}\Omega g \cap m^{-1}SWk_g^{-1}) \cdot \int_{\Lambda \setminus L} f d\mu, \end{aligned}$$

where $k_g \in K$ such that $k_gB = g^{-1}B$.

Let $\Omega_m = g^{-1}\Omega g \cap m^{-1}SWk_g^{-1}$. One can check that $\nu(\partial\Omega_m) = 0$. To show the claim, we first fix $s \in V$ and $\varepsilon > 0$. Note that

$$\Omega_{sm}(a) = \{k \in g^{-1}\Omega g : kk_gB \in a^{-1}m^{-1}(s^{-1}V)WB\}.$$

By Lemma 4.3(1), there exists $C > 0$ such that

$$(6.11) \quad \int_{\Omega_m - \Omega_{sm}(a)} |f(ygk^{-1}a^{-1}(sm)^{-1})| d\nu(k) \leq \varepsilon \quad \text{for all } a \in A_T^+(C).$$

and by Theorem 6.2 for any sufficiently large $C > 0$ and $a \in A_T^+(C)$, we have

$$\left| \int_{\Omega_m} f(ygk^{-1}a^{-1}(sm)^{-1}) d\nu(k) - \nu(\Omega_m) \int_{\Lambda \setminus L} f d\mu \right| < \varepsilon.$$

Hence,

$$\begin{aligned}
& \left| \int_{A_T^+(C)} \int_{\Omega_{sm}(a)} f(ygk^{-1}a^{-1}(sm)^{-1}) \xi(\log a) d\nu(k) da \right. \\
& \quad \left. - \int_{A_T^+(C)} \xi(\log a) da \cdot \nu(\Omega_m) \cdot \int_{\Lambda \setminus L} f d\mu \right| \\
& \leq \int_{A_T^+(C)} \int_{\Omega_m - \Omega_{sm}(a)} |f(ygk^{-1}a^{-1}(sm)^{-1})| \xi(\log a) d\nu(k) da \\
& \quad + \int_{A_T^+(C)} \left| \int_{\Omega_m} f(ygk^{-1}a^{-1}(sm)^{-1}) d\nu(k) - \nu(\Omega_m) \int_{\Lambda \setminus L} f d\mu \right| \xi(\log a) da \\
& \leq 2\varepsilon \int_{A_T^+(C)} \xi(\log a) da \leq 2\varepsilon \int_{A_T^+} \xi(\log a) da.
\end{aligned}$$

Thus, by (6.7) and (6.8),

$$\begin{aligned}
& \left| \int_{A_T^+} \int_{\Omega_{sm}(a)} f(ygk^{-1}a^{-1}(sm)^{-1}) \xi(\log a) d\nu(k) da \right. \\
& \quad \left. - \rho(B_T(e)) \nu(\Omega_m) \int_{\Lambda \setminus L} f d\mu \right| \leq 2\varepsilon \rho(B_T(e)) + o(\rho(B_T(e))).
\end{aligned}$$

This shows that for every $s \in V$,

$$\begin{aligned}
\lim_{T \rightarrow \infty} \frac{1}{\rho(B_T(e))} \int_{A_T^+} \int_{\Omega_{sm}(a)} f(ygk^{-1}a^{-1}(sm)^{-1}) \xi(\log a) d\nu(k) da \\
= \nu(\Omega_m) \int_{\Lambda \setminus L} f d\mu.
\end{aligned}$$

Therefore, (6.10) follows from the Lebesgue dominated convergence theorem.

Step 2: We claim that for every $m \in M$,

(6.12)

$$\lim_{T \rightarrow \infty} \frac{1}{\rho(B_T(e))} \int_{S - \bar{V}} \int_{A_T^+} \int_{\Omega_{sm}(a)} f(ygk^{-1}a^{-1}(sm)^{-1}) \xi(\log a) d\nu(k) dad\sigma(s) = 0.$$

Let $s \in S - \bar{V}$ and $\varepsilon > 0$. By Lemma 4.3(2), there exists $C > 0$ such that

$$\nu(\Omega_{sm}(a)) < \varepsilon \quad \text{for all } a \in A_T^+(C).$$

Hence, by (6.7),

$$\begin{aligned} & \frac{1}{\rho(B_T(e))} \int_{A_T^+(C)} \int_{\Omega_{sm}(a)} |f(ygk^{-1}a^{-1}(sm)^{-1})| \xi(\log a) d\nu(k) da \\ & \leq \frac{\varepsilon \cdot \sup |f|}{\rho(B_T(e))} \int_{A_T^+(C)} \xi(\log a) da \leq \varepsilon \cdot \sup |f|. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, it follows from (6.8) that for any $s \in S - \overline{V}$,

$$\lim_{T \rightarrow \infty} \frac{1}{\rho(B_T(e))} \int_{A_T^+} \int_{\Omega_{sm}(a)} f(ygk^{-1}a^{-1}(sm)^{-1}) \xi(\log a) d\nu(k) da = 0.$$

Hence, (6.12) follows from the Lebesgue dominated convergence theorem.

Since $\sigma(\partial V) = 0$, combining (6.10) and (6.12), we deduce that for every $m \in M$,

$$\begin{aligned} \lim_{T \rightarrow \infty} \frac{1}{\rho(B_T(e))} \int_S \int_{A_T^+} \int_{\Omega_{sm}(a)} f(ygk^{-1}a^{-1}(sm)^{-1}) \xi(\log a) d\nu(k) dad\sigma(s) \\ = \sigma(V) \cdot \nu(g^{-1}\Omega g \cap m^{-1}SWk_g^{-1}) \cdot \int_{\Lambda \setminus L} f d\mu. \end{aligned}$$

Thus, by (4.2), the Lebesgue dominated convergence theorem, and Lemma 4.5,

$$\begin{aligned} \lim_{T \rightarrow \infty} \frac{1}{\rho(B_T(e))} \int_K \int_{A_T^+} \int_{\Omega_{k_1}(a)} f(ygk_2^{-1}a^{-1}(k_1)^{-1}) \xi(\log a) d\nu(k_2) dad\nu(k_1) \\ = \sigma(V) \cdot \int_M \nu(g^{-1}\Omega g \cap m^{-1}SWk_g^{-1}) d\tau(m) \cdot \int_{\Lambda \setminus L} f d\mu \\ = \sigma(V) \nu(g^{-1}\Omega g) \tau(W) \int_{\Lambda \setminus L} f d\mu = \nu(U) \nu_0(\Omega) \int_{\Lambda \setminus L} f d\mu. \end{aligned}$$

Finally, the proposition follows from (6.5) and (6.9). \square

Proof of Theorem 6.1. Let B be a maximal connected split solvable subgroup of G contained in Q . Since $G = KB$, we have $Q = DB$ for $D = K \cap Q$. Then

$$(6.13) \quad Q_T(g, \Omega) = DB_T(g, \Omega)$$

and for suitable Haar measures ρ_B and ν_D on B and D respectively.

$$(6.14) \quad \int_Q f(q) d\rho(q) = \int_D \int_B f(db) d\rho_B(b) d\nu_D(d), \quad f \in C_c(Q).$$

Hence, Theorem 6.1 for Q follows from Theorem 6.1 for B and the Lebesgue dominated convergence theorem. Thus, we may assume that $Q = B$.

Let $\varepsilon > 0$. One can find a neighborhood U of e in K as in Proposition 6.6 and functions $f^+, f^- \in C_c(\Lambda \setminus L)$ such that

$$f^-(xk^{-1}) \leq f(x) \leq f^+(xk^{-1}) \quad \text{for all } x \in \Lambda \setminus L \text{ and } k \in U$$

and

$$\int_{\Lambda \setminus L} |f^+ - f^-| d\mu \leq \varepsilon.$$

Put

$$F_T(k) = \frac{1}{\rho(B_T(g))} \int_{B_T(g, \Omega)} f(yb^{-1}k^{-1}) d\rho(b), \quad k \in K.$$

For any $k \in U$,

$$(6.15) \quad \begin{aligned} \frac{1}{\rho(B_T(g))} \int_{B_T(g, \Omega)} f^-(yb^{-1}k^{-1}) d\rho(b) &\leq F_T(e) \\ &\leq \frac{1}{\rho(B_T(g))} \int_{B_T(g, \Omega)} f^+(yb^{-1}k^{-1}) d\rho(b). \end{aligned}$$

Integrating over $U \subset K$, we obtain

$$(6.16) \quad \begin{aligned} \frac{1}{\rho(B_T(g))} \int_U \int_{B_T(g, \Omega)} f^-(yb^{-1}k^{-1}) d\rho(b) d\nu(k) &\leq \nu(U) \cdot F_T(e) \\ &\leq \frac{1}{\rho(B_T(g))} \int_U \int_{B_T(g, \Omega)} f^+(yb^{-1}k^{-1}) d\rho(b) d\nu(k). \end{aligned}$$

Hence, by Proposition 6.6,

$$\begin{aligned} \nu(U)\nu_0(M_0\Omega) \int_{\Lambda \setminus L} f^- d\mu &\leq \nu(U) \cdot \liminf_{T \rightarrow \infty} F_T(e) \\ &\leq \nu(U) \cdot \limsup_{T \rightarrow \infty} F_T(e) \leq \nu(U)\nu_0(M_0\Omega) \int_{\Lambda \setminus L} f^+ d\mu. \end{aligned}$$

This shows that

$$\begin{aligned} \nu_0(M_0\Omega) \cdot \left(\int_{\Lambda \setminus L} f d\mu - \varepsilon \right) &\leq \liminf_{T \rightarrow \infty} F_T(e) \\ &\leq \limsup_{T \rightarrow \infty} F_T(e) \leq \nu_0(M_0\Omega) \cdot \left(\int_{\Lambda \setminus L} f d\mu + \varepsilon \right), \end{aligned}$$

and the theorem follows. \square

Since it follows from the result of N. Shah (see [S, Theorem 1.1]) and Ratner's topological rigidity [R2] that $\overline{yQ} = \overline{yG_nQ}$ for every $y \in \Lambda \setminus L$, one may expect that Theorem 6.1 holds under the condition $\overline{yQ} = \Lambda \setminus L$ as well.

Lemma 6.17. *Let L be a connected semisimple Lie group with finite center and $L = L_c L_n$ the decomposition of L into the product of compact and noncompact factors. Suppose that for $y \in \Lambda \backslash L$, we have $\overline{yG_n} \supset yL_n$ and $\overline{yG_n Q} = \Lambda \backslash L$. Then the conclusion of Theorem 6.1 holds.*

Proof. Let $B \subset Q$ be a maximal connected split solvable subgroup of G . Then $Q = DB$ for $D = K \cap Q$. By Ratner's theorem on orbit closures [R2], the set $\overline{yG_n}$ is a homogeneous space yG_0 with a probability G_0 -invariant measure μ_0 , where G_0 is a closed connected subgroup of L that contains L_n . Applying Theorem 6.1 to the space yG_0 and the subgroup B , we deduce that

$$\lim_{T \rightarrow \infty} \frac{1}{\rho_B(B_T(g))} \int_{B_T(g, \Omega)} f(yb^{-1}) d\rho_B(b) = \nu_0(M_0 \Omega) \int_{yG_0} f d\mu_0$$

for every $f \in C_c(\Lambda \backslash L)$, where ρ_B is a right invariant Haar measure on B . Since $\overline{yG_n Q} = \Lambda \backslash L$, $L_n Q = L_n D$, and $L_n \subset G_0$, we have

$$(6.18) \quad \Lambda \backslash L = \overline{yG_n D} = yG_0 D.$$

(Here we used that D is compact.) By (6.13), (6.14), and the Lebesgue dominated convergence theorem,

$$\lim_{T \rightarrow \infty} \frac{1}{\rho(Q_T(g))} \int_{Q_T(g, \Omega)} f(yq^{-1}) d\rho(q) = \nu_0(M_0 \Omega) \int_{\Lambda \backslash G} f d\tilde{\mu}$$

for every $f \in C_c(\Lambda \backslash L)$, where the measure $\tilde{\mu}$ is defined by

$$\int_{\Lambda \backslash L} f d\tilde{\mu} = \int_D \int_{yG_0} f(zd^{-1}) d\mu_0(z) d\nu_D(d), \quad f \in C_c(\Lambda \backslash L).$$

Since L_n is a normal subgroup of L , it is clear that the measure $\tilde{\mu}$ is invariant under $L_n D$. This measure corresponds to a Radon measure $\tilde{\mu}$ on L which is right $(L_n D)$ -invariant and left $\text{Stab}_L(y)$ -invariant. Namely,

$$\tilde{\mu}(f) = \int_{\Lambda \backslash L} \left(\sum_{\lambda \in \text{Stab}_L(y)} f(\lambda g) \right) d\tilde{\mu}(g), \quad f \in C_c(L).$$

We have a decomposition $\tilde{\mu} = \int_{L_c} \tilde{\mu}_x d\omega(x)$ for a Radon measure ω on L_c , where $\tilde{\mu}_x$ is a right L_n -invariant measure on the leaf xL_n for ω -a.e. $x \in L_c$. Since L_n commutes with $x \in L_c$, $\tilde{\mu}_x$ is left L_n -invariant too. It follows that the measure $\tilde{\mu}$ is left L_n -invariant. Thus, it is left invariant under $\overline{\text{Stab}_L(y)L_n} \supset G_0$. Setting $E = G_0 \cap L_c$ and $F = DL_n \cap L_c$, we deduce that the measure ω is left E -invariant and right F -invariant. Note that EF is a closed subset of L_c , and it follows from (6.18) that $yL_n EF = \Lambda \backslash L$. Thus, by the Baire category theorem, the set EF contains an open subset of L_c . Since the group $E \times F$ acts transitively on EF , this implies that EF is open in L_c . Thus, $L_c = EF$ because L is connected. By

[K, Theorem 8.32], ω is a Haar measure on L_c . This implies that $\tilde{\mu}$ is L -invariant, and the lemma follows. \square

7. DISTRIBUTION OF LATTICE POINTS IN SECTORS AND THE BOUNDARY

In this section, we apply Theorem 6.1 in the case when $\Lambda \backslash L = \Gamma \backslash G$ in order to deduce Theorems 1.1 and 1.2. We use notations from Theorem 1.2. In particular, A is a split Cartan subgroup with respect to K_1 , $g^{-1}K_1g = K_2$, M_1 is the centralizer of A in K_1 , and $M_2 = K_2 \cap Q$.

To simplify notations, we make the following conventions in this section:

- For $R \subset G$ and $T > 0$,

$$R_T = \{r \in R : d(K_1, K_1gr) < T\}.$$

- For $T, C > 0$, $\Omega_1 \subset K_1$, and $\Omega_2 \subset K_2$,

$$\begin{aligned} N_T(\Omega_1, \Omega_2) &= \#(\Gamma_T \cap g^{-1}K_1A^+\Omega_1 \cap \Omega_2Q), \\ N_T^C(\Omega_1, \Omega_2) &= \#(\Gamma_T \cap g^{-1}K_1A^+(C)\Omega_1 \cap \Omega_2Q), \end{aligned}$$

where $A^+(C)$ is defined as in (5.10).

- For $T, C > 0$ and $\Omega \subset K_1$,

$$\begin{aligned} Q_T(\Omega) &= Q_T \cap g^{-1}K_1A^+\Omega, \\ Q_T^C(\Omega) &= Q_T \cap g^{-1}K_1A^+(C)\Omega. \end{aligned}$$

Let m denote the Haar measure on G such that (5.1) holds for K_1 and ρ the right invariant Haar measure on Q such that

$$(7.1) \quad \int_G f \, dm = \int_{K_2} \int_Q f(kq) \, d\rho(q) \, d\nu_2(k), \quad f \in C_c(G).$$

Lemma 7.2. *For any $C > 0$, $\Omega_1 \subset K_1$, and $\Omega_2 \subset K_2$,*

$$\lim_{T \rightarrow \infty} \frac{1}{m(G_T)} (N_T(\Omega_1, \Omega_2) - N_T^C(\Omega_1, \Omega_2)) = 0.$$

Proof. Fix $D > C > 0$ and $\varepsilon > 0$. By Theorem 3.7, there exists a neighborhood \mathcal{O} of e in G such that

$$(7.3) \quad \mathcal{O}^{-1}g^{-1}K_1A^+(D)K_1 \subset g^{-1}K_1A^+(C)K_1.$$

In addition, we may choose \mathcal{O} so that

$$\Gamma \cap \mathcal{O}^{-1}\mathcal{O} = \{e\} \quad \text{and} \quad \mathcal{O}G_T \subset G_{T+\varepsilon} \quad \text{for all } T > 0.$$

It follows from (7.3) that

$$\mathcal{O} \cdot (\Gamma - g^{-1}K_1A^+(C)K_1) \subset G - g^{-1}K_1A^+(D)K_1.$$

Thus,

$$\begin{aligned}
 N_T(\Omega_1, \Omega_2) - N_T^C(\Omega_1, \Omega_2) &\leq \#\{\gamma \in \Gamma_T - g^{-1}K_1A^+(C)K_1\} \\
 &= \frac{1}{m(\mathcal{O})}m\left(\bigcup_{\gamma \in \Gamma_T - g^{-1}K_1A^+(C)K_1} \mathcal{O}\gamma\right) \leq \frac{1}{m(\mathcal{O})}m(G_{T+\varepsilon} - g^{-1}K_1A^+(D)K_1) \\
 &= \frac{1}{m(\mathcal{O})}m(K_1(A_{T+\varepsilon}^+ - A_{T+\varepsilon}^+(D))K_1) = o(m(G_{T+\varepsilon}))
 \end{aligned}$$

by (5.3) and Lemma 5.11. Now the lemma follows from Lemma 5.9. \square

The proof of the following lemma is similar and is left to the readers.

Lemma 7.4. *For any $C > 0$ and $\Omega \subset K_1$,*

$$\lim_{T \rightarrow \infty} \frac{1}{m(G_T)}(\rho(Q_T(\Omega)) - \rho(Q_T^C(\Omega))) = 0.$$

Proof of Theorem 1.2. Without loss of generality, $m(\Gamma \backslash G) = 1$. It is easy to check that $\nu_1(\partial(M_1\Omega_1)) = \nu_2(\partial(\Omega_2M_2)) = 0$. Thus, we may assume that $\Omega_1 = M_1\Omega_1$ and $\Omega_2 = \Omega_2M_2$.

We need to show that

$$N_T(\Omega_1, \Omega_2) \sim_{T \rightarrow \infty} \nu_1(\Omega_1)\nu_2(\Omega_2)m(G_T).$$

Fix any $\varepsilon > 0$ and $C > 0$. Let U be a neighborhood of e in K_1 with boundary of measure zero such that $\nu_1(\Omega_1^+ - \Omega_1^-) < \varepsilon$, where

$$\Omega_1^+ = \bigcup_{u \in U} u\Omega_1 \quad \text{and} \quad \Omega_1^- = \bigcap_{u \in U} u^{-1}\Omega_1.$$

One can check that $\nu_1(\partial\Omega_1^\pm) = 0$.

By the strong wavefront lemma (Theorem 3.7), there exists a symmetric neighborhood \mathcal{O}' of e in G such that

$$(7.5) \quad \mathcal{O}'g^{-1}K_1A^+(C) \subset g^{-1}K_1A^+U \quad \text{and} \quad \mathcal{O}'G_T \subset G_{T+\varepsilon} \quad \text{for all } T > 0.$$

Set $\mathcal{O} = M_2(\mathcal{O}' \cap Q)M_2$. Note that \mathcal{O} is a symmetric neighborhood of e in Q . Using the fact that $M_2 \subset g^{-1}K_1g$, it is easy to check that (7.5) holds for \mathcal{O} as well. We may also assume that $\rho(\partial\mathcal{O}) = 0$.

Let f be the characteristic function of $\Omega_2\mathcal{O} \subset G$. Since the decomposition $h = h_{K_2}h_Q$ for $h_{K_2} \in K_2$ and $h_Q \in Q$ is uniquely determined modulo M_2 and $M_2\mathcal{O} = \mathcal{O}$, we have

$$f(h) = \chi_{\Omega_2}(h_{K_2})\chi_{\mathcal{O}}(h_Q).$$

We also define a function on $\Gamma \backslash G$ by

$$F(h) = \sum_{\gamma \in \Gamma} f(\gamma h).$$

Step 1: We claim that for any $T > 0$,

$$(7.6) \quad N_T^C(\Omega_1, \Omega_2) \leq \frac{1}{\rho(\mathcal{O})} \int_{Q_{T+\varepsilon}(\Omega_1^+)} F(q^{-1}) d\rho(q),$$

We have

$$(7.7) \quad \begin{aligned} \int_{Q_{T+\varepsilon}(\Omega_1^+)} F(q^{-1}) d\rho(q) &= \int_{Q_{T+\varepsilon}(\Omega_1^+)} \left(\sum_{\gamma \in \Gamma} \chi_{\Omega_2}(\gamma_{K_2}) \chi_{\mathcal{O}}(\gamma_Q q^{-1}) \right) d\rho(q) \\ &= \sum_{\gamma \in \Gamma: \gamma_{K_2} \in \Omega_2} \rho(Q_{T+\varepsilon}(\Omega_1^+) \cap \mathcal{O}\gamma_Q). \end{aligned}$$

It follows from (7.5) that

$$\mathcal{O}Q_T^C(\Omega_1) \subset Q_{T+\varepsilon}(\Omega_1^+).$$

This implies that for every $\gamma \in \Gamma$ such that $\gamma_Q \in Q_T^C(\Omega_1)$,

$$\rho(Q_{T+\varepsilon}(\Omega_1^+) \cap \mathcal{O}\gamma_Q) = \rho(\mathcal{O}).$$

Thus,

$$\begin{aligned} \int_{Q_{T+\varepsilon}(\Omega_1^+)} F(q^{-1}) d\rho(q) &\geq \#\{\gamma \in \Gamma : \gamma_{K_2} \in \Omega_2, \gamma_Q \in Q_T^C(\Omega_1)\} \cdot \rho(\mathcal{O}) \\ &= N_T^C(\Omega_1, \Omega_2) \rho(\mathcal{O}). \end{aligned}$$

This proves (7.6).

Step 2: We claim that for any $T > 0$,

$$(7.8) \quad N_T(\Omega_1, \Omega_2) \geq \frac{1}{\rho(\mathcal{O})} \int_{Q_{T-\varepsilon}^C(\Omega_1^-)} F(q^{-1}) d\rho(q).$$

As in (7.7),

$$(7.9) \quad \int_{Q_{T-\varepsilon}^C(\Omega_1^-)} F(q^{-1}) d\rho(q) = \sum_{\gamma \in \Gamma: \gamma_{K_2} \in \Omega_2} \rho(Q_{T-\varepsilon}^C(\Omega_1^-) \cap \mathcal{O}\gamma_Q).$$

Since $U\Omega_1^- \subset \Omega_1$, we have by (7.5),

$$\mathcal{O}^{-1}Q_{T-\varepsilon}^C(\Omega_1^-) \subset Q_T(\Omega_1).$$

Therefore, for $\gamma \in \Gamma$ such that $\gamma_Q \notin Q_T(\Omega_1)$,

$$\rho(Q_{T-\varepsilon}^C(\Omega_1^-) \cap \mathcal{O}\gamma_Q) = 0.$$

By (7.9),

$$\begin{aligned} \int_{Q_{T-\varepsilon}^C(\Omega_1^-)} F(q^{-1}) d\rho(q) &\leq \#\{\gamma \in \Gamma : \gamma_{K_2} \in \Omega_2, \gamma_Q \in Q_T(\Omega_1)\} \cdot \rho(\mathcal{O}) \\ &= N_T(\Omega_1, \Omega_2) \rho(\mathcal{O}). \end{aligned}$$

This proves (7.8).

Since the boundary of the set $\Omega_2\mathcal{O}$ has measure 0 (this can be checked using (7.1)), the function f can be approximated by continuous functions with compact support and Theorem 6.1 can be applied to the function F (see Lemma 6.17):

$$(7.10) \quad \begin{aligned} \lim_{T \rightarrow \infty} \frac{1}{\rho(Q_T)} \int_{Q_T(\Omega_1^\pm)} F(q^{-1}) d\rho(q) &= \nu_1(\Omega_1^\pm) \int_{G/\Gamma} F dm \\ &= \nu_1(\Omega_1^\pm) \int_G f dm = \nu_1(\Omega_1^\pm) \nu_2(\Omega_2) \rho(\mathcal{O}). \end{aligned}$$

Note that by (7.1), $m(G_T) = \rho(Q_T)$. Combining (7.8), Lemma 7.4, and (7.10), we deduce that

$$\begin{aligned} \liminf_{T \rightarrow \infty} \frac{N_T(\Omega_1, \Omega_2)}{m(G_T)} &\geq \liminf_{T \rightarrow \infty} \frac{1}{m(G_T) \rho(\mathcal{O})} \int_{Q_{T-\varepsilon}^C(\Omega_1^-)} F(q^{-1}) d\rho(q) \\ &\geq \left(\liminf_{T \rightarrow \infty} \frac{m(G_{T-\varepsilon})}{m(G_T)} \right) \cdot \liminf_{T \rightarrow \infty} \frac{1}{\rho(\mathcal{O}) \rho(Q_{T-\varepsilon})} \int_{Q_{T-\varepsilon}(\Omega_1^-)} F(q^{-1}) d\rho(q) \\ &\geq a(\varepsilon) \nu_1(\Omega_1^-) \nu_2(\Omega_2) \geq a(\varepsilon) (\nu_1(\Omega_1) - \varepsilon) \nu_2(\Omega_2), \end{aligned}$$

where $a(\varepsilon)$ is defined in Lemma 5.9. Since $\varepsilon > 0$ is arbitrary, it follows from Lemma 5.9 that

$$\liminf_{T \rightarrow \infty} \frac{N_T(\Omega_1, \Omega_2)}{m(G_T)} \geq \nu_1(\Omega_1) \nu_2(\Omega_2).$$

By Lemma 7.2, (7.6), and (7.10),

$$\begin{aligned} \limsup_{T \rightarrow \infty} \frac{N_T(\Omega_1, \Omega_2)}{m(G_T)} &= \limsup_{T \rightarrow \infty} \frac{N_T^C(\Omega_1, \Omega_2)}{m(G_T)} \\ &\leq \limsup_{T \rightarrow \infty} \frac{1}{m(G_T) \rho(\mathcal{O})} \int_{Q_{T+\varepsilon}(\Omega_1^+)} F(q^{-1}) d\rho(q) \\ &\leq \left(\limsup_{T \rightarrow \infty} \frac{m(G_{T+\varepsilon})}{m(G_T)} \right) \cdot \limsup_{T \rightarrow \infty} \frac{1}{\rho(\mathcal{O}) \rho(Q_{T+\varepsilon})} \int_{Q_{T+\varepsilon}(\Omega_1^+)} F(q^{-1}) d\rho(q) \\ &\leq b(\varepsilon) \nu_1(\Omega_1^+) \nu_2(\Omega_2) \leq b(\varepsilon) (\nu_1(\Omega_1) + \varepsilon) \nu_2(\Omega_2), \end{aligned}$$

where $b(\varepsilon)$ is defined in Lemma 5.9. Thus, by Lemma 5.9,

$$\limsup_{T \rightarrow \infty} \frac{N_T(\Omega_1, \Omega_2)}{m(G_T)} \leq \nu_1(\Omega_1) \nu_2(\Omega_2),$$

and the theorem is proved. \square

Proof of Theorem 1.1. Note that G is a connected semisimple center-free Lie group with no compact factors, and K_x and K_y are maximal compact subgroups of G . Since G acts transitively on X (see, for example, [H, Theorem IV.3.3]), there exists $g \in G$ such that $y = xg$. Then $K_y = g^{-1}K_xg$. The closed positive Weyl

chamber \mathcal{W}_x at x is of the form xA^+ , where A^+ is a positive Weyl chamber in a split Cartan subgroup A of G with respect to K_x . The stabilizer of $b \in X(\infty)$ in G is a parabolic subgroup Q of G (see [GJT, Proposition III.3.8]). In particular, Q contains a maximal connected split solvable subgroup of G . Let $\pi : K_y \rightarrow bG$ be the map given by $k \mapsto bk^{-1}$. Then for $\Omega_1 \subset K_x$ and $\Omega_2 \subset bG$,

$$\begin{aligned} & \#\{\gamma \in \Gamma : y\gamma \in \mathcal{W}_x\Omega_1 \cap B_T(x), b\gamma^{-1} \in \Omega_2\} \\ &= \#\{\gamma \in \Gamma \cap g^{-1}K_xA^+\Omega_1 \cap \pi^{-1}(\Omega_2)Q : d(K_x, K_xg\gamma) < T\}. \end{aligned}$$

Note that π maps the probability Haar measure on K_y to the measure $m_{b,y}$. Using the fact that the map π is open, one can check that the set $\pi^{-1}(\Omega_2)$ has boundary of measure zero if $m_{b,y}(\partial\Omega_2) = 0$. Hence, Theorem 1.1 follows from Theorem 1.2. \square

8. DISTRIBUTION OF LATTICE POINTS IN BISECTORS

Let G be a connected semisimple Lie group with finite center and $G = KA^+K$ a Cartan decomposition of G . To simplify notations, we fix $g \in G$ and for $\Omega_1, \Omega_2 \subset K$ and $T, C > 0$, define

$$\begin{aligned} G_T &= \{h \in G : d(K, Kgh) < T\}, \\ G_T(\Omega_1, \Omega_2) &= g^{-1}\Omega_1A^+\Omega_2 \cap G_T, \\ N_T(\Omega_1, \Omega_2) &= \#(\Gamma \cap G_T(\Omega_1, \Omega_2)). \end{aligned}$$

If we set $A_T^+ = \{a \in A^+ : d(K, Ka) < T\}$, then $G_T(\Omega_1, \Omega_2) = g^{-1}\Omega_1A_T^+\Omega_2$.

Let G be a closed subgroup of a Lie group L and Λ a lattice in L . Let m be a Haar measure on G such that (5.1) holds and μ the Haar measure on L such that $\mu(\Lambda \backslash L) = 1$.

Theorem 8.1. *Suppose that for $y \in \Lambda \backslash L$, the orbit yG_n is dense in $\Lambda \backslash L$, where G_n is the product of all noncompact simple factors of G . For any Borel subsets $\Omega_1, \Omega_2 \subset K$ with boundaries of measure zero and $f \in C_c(\Lambda \backslash L)$,*

$$\lim_{T \rightarrow \infty} \frac{1}{m(G_T)} \int_{G_T(\Omega_1, \Omega_2)} f(yh^{-1}) dm(h) = \nu(\Omega_1)\nu(\Omega_2) \int_{\Lambda \backslash L} f d\mu.$$

Proof. By (5.1),

$$\begin{aligned} (8.2) \quad \int_{G_T(\Omega_1, \Omega_2)} f(yh^{-1}) dm(h) &= \int_{\Omega_1} \int_{A_T^+} \int_{\Omega_2} f(yk_2^{-1}a^{-1}k_1^{-1}g) \xi(\log a) dk_2 dadk_1 \\ &= \int_{\Omega_1} \int_{A_T^+} \int_{\Omega_2^{-1}} f(yk_2a^{-1}k_1^{-1}g) \xi(\log a) dk_2 dadk_1. \end{aligned}$$

Since $\nu(\Omega_2^{-1}) = \nu(\Omega_2)$, by Theorem 6.2, for every $\varepsilon > 0$, there exists $C > 0$ such that

$$(8.3) \quad \left| \int_{\Omega_2^{-1}} f(yk_2a^{-1}k_1^{-1}g) dk_2 - \nu(\Omega_2) \int_{\Lambda \backslash L} f d\mu \right| < \varepsilon$$

for all $a \in A_T^+(C)$. Since $m(G_T) = \int_{A_T^+} \xi(\log a) da$, it follows from Lemma 5.11 that

$$(8.4) \quad \int_{A_T^+ - A_T^+(C)} \xi(\log a) da = o(m(G_T)) \quad \text{as } T \rightarrow \infty.$$

Combining (8.3) and (8.4), we get

$$\begin{aligned} & \left| \int_{A_T^+} \int_{\Omega_2^{-1}} f(yk_2a^{-1}k_1^{-1}g) dk_2 \xi(\log a) da - \nu(\Omega_2) \cdot \int_{A_T^+} \xi(\log a) da \cdot \int_{\Lambda \backslash L} f d\mu \right| \\ & \leq \int_{A_T^+(C)} \left| \int_{\Omega_2^{-1}} f(yk_2a^{-1}k_1^{-1}g) dk_2 - \nu(\Omega_2) \int_{\Lambda \backslash L} f d\mu \right| \xi(\log a) da \\ & + 2 \sup |f| \int_{A_T^+ - A_T^+(C)} \xi(\log a) da \leq \varepsilon \int_{A_T^+(C)} \xi(\log a) da + o(m(G_T)) \\ & \leq \varepsilon \cdot m(G_T) + o(m(G_T)) \quad \text{as } T \rightarrow \infty. \end{aligned}$$

This proves that for every $k_1 \in K$,

$$\lim_{T \rightarrow \infty} \frac{1}{m(G_T)} \int_{A_T^+} \int_{\Omega_2^{-1}} f(yk_2a^{-1}k_1^{-1}g) \xi(\log a) dk_2 da = \nu(\Omega_2) \int_{\Lambda \backslash L} f d\mu.$$

Now the statement follows from (8.2) and the Lebesgue dominated convergence theorem. \square

Proof of Theorem 1.6. We need to show that

$$N_T(\Omega_1 M, M \Omega_2) \sim_{T \rightarrow \infty} \nu(\Omega_1 M) \nu(M \Omega_2) \cdot \frac{m(G_T)}{m(\Gamma \backslash G)}.$$

We may assume without loss of generality that $\Omega_1 M = \Omega_1$, $M \Omega_2 = \Omega_2$, and $m(\Gamma \backslash G) = 1$. Since M contains all compact factors of G , we may assume that G contains no compact factors.

Fix $\varepsilon > 0$ and $C > 0$. There exists a neighborhood U of e in K with boundary of measure zero such that $\nu(\Omega_i^+ - \Omega_i^-) < \varepsilon$, $i = 1, 2$, where

$$\Omega_i^+ = \bigcup_{u \in U} \Omega_i u \quad \text{and} \quad \Omega_i^- = \bigcap_{u \in U} \Omega_i u^{-1}.$$

Note that $\nu(\partial\Omega_i^\pm) = 0$, $\Omega_i U \subset \Omega_i^+$, and $\Omega_i^- U \subset \Omega_i^-$. By the strong wavefront lemma (Theorem 3.7), there exists a neighborhood \mathcal{O} of e in G such that

$$(8.5) \quad \begin{aligned} \mathcal{O}^{-1}\Omega_1 A^+(C)\Omega_2 &\subset \Omega_1^+ A^+ \Omega_2^+, \\ \mathcal{O}\Omega_1^- A^+(C)\Omega_2^- &\subset \Omega_1^- A^+ \Omega_2^-, \\ \mathcal{O}^{\pm 1}G_T &\subset G_{T+\varepsilon} \quad \text{for all } T > 0. \end{aligned}$$

Let

$$\begin{aligned} G_T^C(\Omega_1, \Omega_2) &= g^{-1}KA^+(C)K \cap G_T(\Omega_1, \Omega_2), \\ N_T^C(\Omega_1, \Omega_2) &= \#(\Gamma \cap G_T^C(\Omega_1, \Omega_2)). \end{aligned}$$

It is not hard to show (see Lemmas 7.2 and 7.4 for a similar argument) that

$$(8.6) \quad m(G_T(\Omega_1, \Omega_2) - G_T^C(\Omega_1, \Omega_2)) = o(m(G_T)),$$

$$(8.7) \quad N_T(\Omega_1, \Omega_2) - N_T^C(\Omega_1, \Omega_2) = o(m(G_T))$$

as $T \rightarrow \infty$.

Let $f \in C_c(G)$ be such that $f \geq 0$, $\text{supp}(f) \subset \mathcal{O}$ and $\int_G f \, dm = 1$. Define a function on $\Gamma \backslash G$ by

$$F(h) = \sum_{\gamma \in \Gamma} f(\gamma h).$$

Clearly, $\int_{\Gamma \backslash G} F \, dm = 1$. We claim that

$$(8.8) \quad N_T^C(\Omega_1, \Omega_2) \leq \int_{G_{T+\varepsilon}(\Omega_1^+, \Omega_2^+)} F(h^{-1}) \, dm(h)$$

and

$$(8.9) \quad N_T(\Omega_1, \Omega_2) \geq \int_{G_{T-\varepsilon}^C(\Omega_1^-, \Omega_2^-)} F(h^{-1}) \, dm(h).$$

First, we observe that

$$\int_{G_{T+\varepsilon}(\Omega_1^+, \Omega_2^+)} F(h^{-1}) \, dm(h) = \sum_{\gamma \in \Gamma} \int_{G_{T+\varepsilon}(\Omega_1^+, \Omega_2^+) \gamma^{-1}} f(h^{-1}) \, dm(h).$$

By (8.5),

$$\mathcal{O}^{-1}G_T^C(\Omega_1, \Omega_2) \subset G_{T+\varepsilon}(\Omega_1^+, \Omega_2^+).$$

Thus, for every $\gamma \in \Gamma \cap G_T^C(\Omega_1, \Omega_2)$, we have $\mathcal{O}^{-1} \subset G_{T+\varepsilon}(\Omega_1^+, \Omega_2^+) \gamma^{-1}$ and

$$\int_{G_{T+\varepsilon}(\Omega_1^+, \Omega_2^+) \gamma^{-1}} f(h^{-1}) \, dm(h) = 1.$$

This implies (8.8).

To prove (8.9), we use that

$$\int_{G_{T-\varepsilon}^C(\Omega_1^-, \Omega_2^-)} F(h^{-1}) \, dm(h) = \sum_{\gamma \in \Gamma} \int_{G_{T-\varepsilon}^C(\Omega_1^-, \Omega_2^-) \gamma^{-1}} f(h^{-1}) \, dm(h).$$

By (8.5),

$$\mathcal{O}G_{T-\varepsilon}^C(\Omega_1^-, \Omega_2^-) \subset G_T(\Omega_1, \Omega_2).$$

Therefore, for $\gamma \in \Gamma - G_T(\Omega_1, \Omega_2)$, we have $\mathcal{O}^{-1} \cap G_{T-\varepsilon}^C(\Omega_1^-, \Omega_2^-)\gamma^{-1} = \emptyset$ and

$$\int_{G_{T-\varepsilon}^C(\Omega_1^-, \Omega_2^-)\gamma^{-1}} f(h^{-1}) dm(h) = 0.$$

This proves (8.9).

By (8.9), (8.6), and Theorem 8.1,

$$\begin{aligned} \liminf_{T \rightarrow \infty} \frac{N_T(\Omega_1, \Omega_2)}{m(G_T)} &\geq \liminf_{T \rightarrow \infty} \frac{1}{m(G_T)} \int_{G_{T-\varepsilon}^C(\Omega_1^-, \Omega_2^-)} F(h^{-1}) dm(h) \\ &\geq \left(\liminf_{T \rightarrow \infty} \frac{m(G_{T-\varepsilon})}{m(G_T)} \right) \cdot \liminf_{T \rightarrow \infty} \frac{1}{m(G_{T-\varepsilon})} \int_{G_{T-\varepsilon}^C(\Omega_1^-, \Omega_2^-)} F(h^{-1}) dm(h) \\ &\geq a(\varepsilon)\nu(\Omega_1^-)\nu(\Omega_2^-) \geq a(\varepsilon)(\nu(\Omega_1) - \varepsilon)(\nu(\Omega_2) - \varepsilon), \end{aligned}$$

where $a(\varepsilon)$ is defined in Lemma 5.9. Since $\varepsilon > 0$ is arbitrary, it follows from Lemma 5.9 that

$$\liminf_{T \rightarrow \infty} \frac{N_T(\Omega_1, \Omega_2)}{m(G_T)} \geq \nu(\Omega_1)\nu(\Omega_2).$$

The opposite inequality for lim sup is proved similarly using (8.7), (8.8), and Theorem 8.1. \square

9. MEASURE-PRESERVING LATTICE ACTIONS

Let G be a connected semisimple Lie group with finite center and Γ_1, Γ_2 lattices in G . Let $L = G \times G$, $\Lambda = \Gamma_1 \times \Gamma_2$, $H = \{(g, g) : g \in G\}$, and Q a closed subgroup of H containing a maximal connected split solvable subgroup of H . Fix an invariant Riemannian metric d on the symmetric space $K \backslash G$, and for $T > 0$ and $g \in G$, define

$$Q_T(g) = \{(q, q) \in Q : d(K, Kgq) < T\}.$$

Corollary 9.1. *Suppose that for $y \in \Lambda \backslash L$, the orbit yQ is dense in $\Lambda \backslash L$. Then for any $f \in C_c(\Lambda \backslash L)$,*

$$\lim_{T \rightarrow \infty} \frac{1}{\rho(Q_T(g))} \int_{Q_T(g)} f(yq^{-1}) d\rho(q) = \int_{\Lambda \backslash L} f d\mu,$$

where ρ is a right Haar measure on Q , and μ is the L -invariant probability measure on $\Lambda \backslash L$.

Proof. It suffices to check the conditions of Lemma 6.17. Namely, we need to show that

$$(9.2) \quad \overline{yH_n} \supset yL_n,$$

where H_n and L_n denote the product of all noncompact simple factors of H and L respectively. We also denote by H_c and L_c the product of all compact simple factors of H and L . By Ratner's theorem on orbit closures [R2], $\overline{yH_n} = yH_0$ for some closed subgroup H_0 of L containing H_n . Then $yH_0H_c = \overline{yH} = \Lambda \backslash L$, and it follows from the Baire category theorem (cf. proof of Lemma 6.17) that H_0H_c is an open subset of L . Since H_0H_c is also closed, we conclude that $L = H_0H_c$. Now (9.2) follows from the Lemma 9.3 below. \square

Lemma 9.3. *Let S be a connected subgroup of L that contains H_n . Then $S = (S \cap L_n)(S \cap L_c)$.*

Proof. Let $\mathfrak{h}_n \subset \mathfrak{s} \subset \mathfrak{l} = \mathfrak{l}_n \oplus \mathfrak{l}_c$ be the corresponding Lie algebras. We have decompositions

$$\mathfrak{h}_n = \bigoplus_i \mathfrak{h}_n^i \quad \text{and} \quad \mathfrak{l}_n = \bigoplus_i \mathfrak{l}_n^i,$$

where \mathfrak{h}_n^i and \mathfrak{l}_n^i are the simple ideals of \mathfrak{h}_n and \mathfrak{l}_n respectively such that $\mathfrak{h}_n^i \subset \mathfrak{l}_n^i$. Note that \mathfrak{h}_n^i is a maximal subalgebra of \mathfrak{l}_n^i . In particular, \mathfrak{h}_n^i is its own normalizer in \mathfrak{l}_n^i .

It suffices to show that for every $s = (\sum_i s_i) + s_c \in \mathfrak{s}$ with $s_i \in \mathfrak{l}_n^i$ and $s_c \in \mathfrak{l}_c$, we have $s_i, s_c \in \mathfrak{s}$. Clearly,

$$[\mathfrak{h}_n^i, s] = [\mathfrak{h}_n^i, s_i] \subset \mathfrak{s}.$$

If $[\mathfrak{h}_n^i, s_i] + \mathfrak{h}_n^i \neq \mathfrak{h}_n^i$, then $[\mathfrak{h}_n^i, s_i] + \mathfrak{h}_n^i$ generates \mathfrak{l}_n^i , and hence, $\mathfrak{l}_n^i \subset \mathfrak{s}$. Otherwise, s_i normalizes \mathfrak{h}_n^i , and it follows that $s_i \in \mathfrak{h}_n^i$. In both cases, $s_i \in \mathfrak{s}$. Then $s_c = s - \sum_i s_i \in \mathfrak{s}$ too. This proves the lemma. \square

Theorem 1.9 can be deduced from Corollary 9.1, as it was explained in [GW] or [O].

REFERENCES

- [DM] S. G. Dani and G. A. Margulis. *Asymptotic behavior of trajectories of unipotent flows on homogeneous spaces*, Proc. Indian Acad. Sci. Math. Sci. 101 (1991), no. 1, 1–17.
- [DRS] W. Duke, Z. Rudnick and P. Sarnak. *Density of integer points on affine homogeneous varieties*, Duke Math. J. 71 (1993), no. 1, 143–179.
- [EMM] A. Eskin, G. Margulis and S. Mozes. *Upper bounds and asymptotics in a quantitative version of the Oppenheim conjecture*, Ann. of Math. (2) 147 (1998), no. 1, 93–141.
- [EM] A. Eskin and C. McMullen. *Mixing, counting and equidistribution on Lie groups*, Duke Math. J. 71 (1993), no. 1, 181–209.
- [EMS] A. Eskin, S. Mozes and N. Shah. *Unipotent flows and counting lattice points on homogeneous varieties*, Ann. of Math. (2) 143 (1996), no. 2, 253–299.
- [EO] A. Eskin and H. Oh. *Integer points on a family of homogeneous varieties and unipotent flows*, To appear in Duke Math. J.
- [H] S. Helgason. *Differential geometry, Lie groups, and symmetric spaces*, Pure and Applied Mathematics, 80. Academic Press, Inc., New York-London, 1978.

- [GO] W. T. Gan and H. Oh. *Equidistribution of integer points on a family of homogeneous varieties: a problem of Linnik*, Compositio Math. 136 (2003), no. 3, 323–352.
- [G] A. Good. *Local analysis of Selberg’s trace formula*, Lecture Notes in Mathematics, 1040, Springer-Verlag, Berlin, 1983.
- [Go] A. Gorodnik. *Lattice action on the boundary of $SL_n(\mathbb{R})$* , Ergodic Theory Dynam. Systems 23 (2003), no. 6, 1817–1837.
- [GM] A. Gorodnik and F. Maucourant. *Proximity and equidistribution on the Furstenberg boundary*, Geom. Dedicata 113 (2005), 197–213.
- [GOS] A. Gorodnik, H. Oh and N. Shah. In preparation.
- [GW] A. Gorodnik and B. Weiss, *Distribution of lattice orbits on homogeneous varieties*, To appear in GAFA.
- [GJT] Y. Guivarc’h, L. Ji and J. C. Taylor. *Compactifications of symmetric spaces*, Progress in Mathematics, 156. Birkhäuser Boston, Inc., Boston, MA, 1998.
- [K] A. Knapp. *Representation theory of semisimple Lie groups, An overview based on examples*, Princeton university press, 1986.
- [Kn] G. Knieper. *On the asymptotic geometry of nonpositively curved manifolds*, Geom. Funct. Anal. 7 (1997), no. 4, 755–782.
- [L] F. Ledrappier. *Ergodic properties of some linear actions*, J. Math. Sci. (New York) 105 (2001), no. 2, 1861–1875.
- [M] G. Margulis. *On some aspects of the theory of Anosov systems*, Springer-Verlag, Berlin, 2004.
- [Ma] F. Maucourant. *Approximation diophantienne, dynamique des chambres de Weyls et répartition d’orbites de réseaux*, PhD Thesis, Université de Lille, 2002.
- [Mor] D. Morris Witte. *Cocompact subgroups of semisimple Lie groups*, Lie algebra and related topics (Madison, WI, 1988), 309–313, Contemp. Math., 110, Amer. Math. Soc., Providence, RI, 1990.
- [Mo] G. D. Mostow. *Strong rigidity of locally symmetric spaces*, Annals of Mathematics Studies, No. 78. Princeton University Press, 1973.
- [N] P. Nicholls. *A lattice point problem in hyperbolic space*, Michigan Math. J. 30 (1983), no. 3, 273–287.
- [No] A. Nogueira. *Orbit distribution on \mathbb{R}^2 under the natural action of $SL(2, \mathbb{Z})$* , Preprint, 2000.
- [O] H. Oh. *Lattice action on finite volume homogeneous spaces*, J. Korean Math. Soc. 42 (2005), no. 4, 635–653.
- [R1] M. Ratner. *On Raghunathan’s measure conjecture*, Ann. of Math. (2) 134 (1991), no. 3, 545–607.
- [R2] M. Ratner. *On Raghunathan’s topological conjecture and distribution of unipotent flows*, Duke Math. J. 63 (1991), no. 1, 235–280.
- [S] N. Shah. *Limit distribution of expanding translates of certain orbits on homogeneous spaces on homogeneous spaces*, Proc. Indian Acad. Sci. Math. Sci. 106 (1996), no. 2, 105–125.
- [Sh] R. Sharp. *Sector estimates for Kleinian groups*, Port. Math. (N.S.) 58 (2001), no. 4, 461–471.
- [V] N. Vatsal. *Uniform distribution of Heegner points*, Invent. Math. 148 (2002), no. 1, 1–46.
- [W] G. Warner. *Harmonic Analysis on Semi-simple Lie groups I*, Springer-Verlag, New York-Heidelberg, 1972.
- [Wi] D. Widder. *The Laplace Transform*, Princeton Mathematical Series, v. 6. Princeton University Press, Princeton, N. J., 1941.
- [Z] R. Zimmer. *Ergodic theory and semisimple groups*, Monographs in Mathematics, 81. Birkhäuser Verlag, Basel, 1984.

MATHEMATICS 253-37, CALTECH, PASADENA, CA 91106

E-mail address: gorodnik@caltech.edu

MATHEMATICS 253-37, CALTECH, PASADENA, CA 91106

E-mail address: heeoh@caltech.edu

Current address: Math Department, 151 Thayer St., Brown University, Providence, RI 02912

E-mail address: heeoh@math.brown.edu