

LIMITS OF TRANSLATES OF DIVERGENT GEODESICS AND INTEGRAL POINTS ON ONE-SHEETED HYPERBOLOIDS

HEE OH AND NIMISH A. SHAH

ABSTRACT. For any non-uniform lattice Γ in $\mathrm{SL}_2(\mathbb{R})$, we describe the limit distribution of orthogonal translates of a *divergent* geodesic in $\Gamma \backslash \mathrm{SL}_2(\mathbb{R})$. As an application, for a quadratic form Q of signature $(2, 1)$, a lattice Γ in its isometry group, and $v_0 \in \mathbb{R}^3$ with $Q(v_0) > 0$, we compute the asymptotic (with a logarithmic error term) of the number of points in a discrete orbit $v_0\Gamma$ of norm at most T , when the stabilizer of v_0 in Γ is finite. Our result in particular implies that for any non-zero integer d , the smoothed count for number of integral binary quadratic forms with discriminant d^2 and with coefficients bounded by T is asymptotic to $c \cdot T \log T + O(T)$.

1. INTRODUCTION

1.1. Motivation. Let $Q \in \mathbb{Z}[x_1, \dots, x_n]$ be a homogeneous polynomial and set $V_m := \{x \in \mathbb{R}^n : Q(x) = m\}$ for an integer m . It is a fundamental problem to understand the set $V_m(\mathbb{Z}) = \{x \in \mathbb{Z}^n : Q(x) = m\}$ of integral solutions.

In particular, we are interested in the asymptotic of the number $N(T) := \#\{x \in V_m(\mathbb{Z}) : \|x\| < T\}$ as $T \rightarrow \infty$, where $\|\cdot\|$ is a fixed norm on \mathbb{R}^n .

The answer to this question depends quite heavily on the geometry of the ambient space V_m . We suppose that the variety V_m is homogeneous, i.e., there exist a connected semisimple real algebraic group G defined over \mathbb{Q} and a \mathbb{Q} -rational representation $\iota : G \rightarrow \mathrm{SL}_n$ such that $V_m = v_0 \cdot \iota(G)$ for some non-zero $v_0 \in \mathbb{Q}^n$.

Let $\Gamma < G(\mathbb{Q})$ be a subgroup commensurable with $G(\mathbb{Z})$ preserving $V_m(\mathbb{Z})$. Since V_m is Zariski closed, by Borel and Harish-Chandra [3, Theorems 6.9 and 7.8], the co-volume of Γ in G is finite and there are only finitely many Γ -orbits in $V_m(\mathbb{Z})$. Hence understanding the asymptotic of $N(T)$ is reduced to the orbital counting problem of estimating $\#(v_0\Gamma \cap B_T)$ for $B_T = \{x \in V_m : \|x\| < T\}$ and $v_0 \in V_m(\mathbb{Z})$.

1991 *Mathematics Subject Classification.* 11N45, 37A17, 22E40 (Primary).

Key words and phrases. Counting lattice points, mixing of geodesic flow, decay of matrix coefficients.

Oh is partially supported by NSF grants 0629322
Shah is partially supported by NSF grant 1001654.

Theorem 1.1 (Duke-Rudnick-Sarnak [10]). *Set H to be the stabilizer subgroup of v_0 in G . Suppose that H is a symmetric subgroup of G . If the volume of $(H \cap \Gamma) \backslash H$ is finite, i.e., if $H \cap \Gamma$ is a lattice in H , we have, as $T \rightarrow \infty$,*

$$\#(v_0\Gamma \cap B_T) \sim \frac{\text{vol}_H(H \cap \Gamma \backslash H)}{\text{vol}_G(\Gamma \backslash G)} \text{vol}_{H \backslash G}(B_T),$$

that is, the ratio of both the sides converges to 1 as $T \rightarrow \infty$, where the volumes on H , G and $v_0G \simeq H \backslash G$ are computed with respect to invariant measures chosen compatibly; that is, $d\text{vol}_G = d\text{vol}_H \times d\text{vol}_{H \backslash G}$ locally.

A simpler proof of this result using mixing of geodesic flow was given by Eskin and McMullen [11]. When H is any maximal reductive \mathbb{Q} -subgroup with $\text{vol}((H \cap \Gamma) \backslash H) < \infty$, the same conclusion was obtained by Eskin, Mozes and Shah in [12] using Ratner's description [17] of measures invariant under unipotent flows. An effective version of Theorem 1.1 has also been obtained in [10] for $(H \cap \Gamma) \backslash H$ compact and in [2] in general.

In view of the the main term of the asymptotic, it is crucial to assume that $\text{vol}(H \cap \Gamma \backslash H) < \infty$ in Theorem 1.1. The main aim of this paper is to break this barrier; to investigate the counting problem in the case when $\text{vol}(H \cap \Gamma \backslash H) = \infty$.

If H is a semisimple Lie group with no compact factors in a semisimple Lie group G , then any closed $\Gamma \backslash \Gamma H$ in $\Gamma \backslash G$ must be of finite volume by Dani [6] and Margulis [16] (see also [20]).

If Q is a quadratic form of signature (p, q) with $p + q \geq 3$, $p \geq q$ and G is the special orthogonal group of Q , the case of $\text{vol}(H \cap \Gamma \backslash H) = \infty$ for $H = \text{Stab}_G(v_0)$ arises only when $(p, q) = (2, 1)$ and $Q(v_0) = m > 0$, that is, when the variety $V_m = \{x \in \mathbb{R}^3 : Q(x) = m\}$ is a one-sheeted hyperboloid. To prove this claim, note that any non-compact stabilizer H of $v_0 \in \mathbb{R}^n$ in G is either locally isomorphic to $\text{SO}(p-1, q)$, $\text{SO}(p, q-1)$ or a compact extension of a horospherical subgroup. Except for the case of $\text{SO}(1, 1)$ the groups $\text{SO}(p-1, q)$ and $\text{SO}(p, q-1)$ are compact or semi-simple. Also any closed orbit of a compact extension of a horospherical subgroup in $\Gamma \backslash G$ is compact (cf. [8]). Therefore in view the above remarks it follows that the case of $\text{vol}(H \cap \Gamma \backslash H) = \infty$ arises only when $H \simeq \text{SO}(1, 1)$; hence $n = 3$ and $Q(v_0) > 0$.

In the next subsection, we state our main theorem in a greater generality, not necessarily in the arithmetic situation.

1.2. Counting integral points on a one-sheeted hyperboloid. Let $Q(x_1, x_2, x_3)$ be an real quadratic form of signature $(2, 1)$. Denote by G the identity component of the special orthogonal group $\text{SO}_Q(\mathbb{R})$. Let $\Gamma < G$ be a lattice and $v_0 \in \mathbb{R}^3$ be such that $Q(v_0) > 0$ and the orbit $v_0\Gamma$ is discrete. As before, we fix a norm $\|\cdot\|$ on \mathbb{R}^3 and set $B_T := \{x \in v_0G : \|w\| < T\}$.

To present our theorem with a best possible error term, we consider the following smoothed counting function: fixing a non-negative function $\psi \in$

$C_c^\infty(G)$ with integral one, let

$$\tilde{N}_T := \sum_{v \in v_0\Gamma} (\chi_{B_T} * \psi)(v)$$

where $\chi_{B_T} * \psi(x) = \int_G \chi_{B_T}(xg)\psi(g) dg$, $x \in v_0G$, is the convolution of the characteristic function of B_T and ψ . Note that $\tilde{N}_T \asymp \#(v_0\Gamma \cap B_T)$ in the sense that their ratio is in between two uniform constants for all sufficiently large $T \gg 1$.

Denoting by $H \simeq \mathrm{SO}(1, 1)^\circ$ the identity component of the one-dimensional stabilizer subgroup of v_0 in G , note that $\mathrm{vol}(H \cap \Gamma \backslash H) < \infty$ if and only if $H \cap \Gamma$ is infinite. In order to state our theorem, we write H as a one-parameter subgroup $\{h(s) : s \in \mathbb{R}\}$ so that the Lebesgue measure ds defines a Haar measure on H : $\int_{-\log T}^{\log T} ds = \mathrm{vol}_H(\{h(s) : |s| < \log T\})$.

Theorem 1.2. *If the volume of $(H \cap \Gamma) \backslash H$ is infinite, we have the following:*

(1) For $T \gg 1$,

$$N_T = \frac{2 \log T \cdot \mathrm{vol}_{H \backslash G}(B_T)}{\mathrm{vol}_G(\Gamma \backslash G)} (1 + O((\log T)^{-0.25})),$$

where $d \mathrm{vol}_G = ds \times d \mathrm{vol}_{H \backslash G}$ locally.

(2) For $T \gg 1$,

$$\tilde{N}_T = c \cdot T \log T + O(T),$$

where $c = \lim_{T \rightarrow \infty} \frac{2 \mathrm{vol}_{H \backslash G}(B_T)}{T \mathrm{vol}_G(\Gamma \backslash G)}$.

We note that when $\mathrm{vol}(H \cap \Gamma \backslash H) < \infty$, $\tilde{N}_T = c \cdot T + O(T^\alpha)$ for $0 < \alpha < 1$ is obtained in [10]. We believe, as suggested by Z. Rudnick to us, that $\tilde{N}_T = c \cdot T \log T + c' \cdot T + O(T^\alpha)$ for some $c' > 0$ and $0 < \alpha < 1$ and hence the order of the second term for \tilde{N}_T cannot be improved.

Theorem 1.2 can be generalized to the orbital counting for more general representations of $\mathrm{SL}_2(\mathbb{R})$ (see section 6).

Remark 1.3. In the case when $Q = x_1^2 + x_2^2 - d^2 x_3^2$ for $d \in \mathbb{Z}$, $v_0 = (1, 0, 0)$, and $\Gamma = \mathrm{SO}_Q(\mathbb{Z})$, it was pointed out in [10] that an elementary number theoretic computation of [19] leads to the asymptotic

$$\#\{(x_1, x_2, x_3) \in v_0\Gamma : \sqrt{x_1^2 + x_2^2 + d^2 x_3^2} < T\} = c \cdot T \log T + O(T \log(\log T)).$$

However this deduction seems to work only for this very special case; for instance, we are not aware of any other approach than ours which can deal with non-arithmetic situations.

1.3. Arithmetic case and Integral binary quadratic forms. In the arithmetic case, Theorem 1.2 (for $\mathrm{vol}(H \cap \Gamma \backslash H) = \infty$) together with Theorem 1.1 (for $\mathrm{vol}(H \cap \Gamma \backslash H) < \infty$) implies the following:

Corollary 1.4. *Let $Q(x_1, x_2, x_3)$ be an integral quadratic form with signature $(2, 1)$. Suppose that for some $v_0 \in \mathbb{Z}^3$ with $Q(v_0) > 0$, the stabilizer subgroup of v_0 is isotropic over \mathbb{Q} .*

Then there exists $c = c(\|\cdot\|) > 0$ such that as $T \rightarrow \infty$,

$$\#\{x \in \mathbb{Z}^3 : Q(x) = Q(v_0), \|x\| < T\} = c \cdot T \log T + O(T(\log T)^{3/4}).$$

For a binary quadratic form $q(x, y) = ax^2 + bxy + cy^2$, its discriminant $\text{disc}(q)$ is defined to be $b^2 - 4ac$. The group $\text{SL}_2(\mathbb{R})$ acts on the space of binary quadratic forms by $(g \cdot q)(x, y) = q(g^{-1}(x, y))$ and preserves the discriminant. For $d \in \mathbb{Z}$, denote by $\mathcal{B}_d(\mathbb{Z})$ the space of integral binary quadratic forms with discriminant d . Note that $\mathcal{B}_d(\mathbb{Z}) \neq \emptyset$ if and only if d congruent to 0 or 1 mod 4. Now d is a square *if and only if* the stabilizer of every $q \in \mathcal{B}_d(\mathbb{Z})$ in $\text{SL}_2(\mathbb{Z})$ is infinite *if and only if* every $q \in \mathcal{B}_d(\mathbb{Z})$ is decomposable over \mathbb{Z} . (cf. [4]).

Therefore Corollary 1.4 implies the following:

Theorem 1.5. *For any non-zero square $d \in \mathbb{Z}$, there exists $c_0 > 0$ such that*

$$\#\{q \in \mathcal{B}_d(\mathbb{Z}) : \text{disc}(q) = d, \|q\| < T\} = c_0 \cdot T \log T + O(T(\log T)^{3/4})$$

where $\|ax^2 + bxy + cy^2\| = \|(a, b, c)\|$.

1.4. Orthogonal translates of a divergent geodesic. Let $G = \text{SL}_2(\mathbb{R})$ and Γ be a non-uniform lattice in G . For $s \in \mathbb{R}$, define

$$h(s) = \begin{bmatrix} \cosh(s/2) & \sinh(s/2) \\ \sinh(s/2) & \cosh(s/2) \end{bmatrix}, \quad a(s) = \begin{bmatrix} e^{s/2} & 0 \\ 0 & e^{-s/2} \end{bmatrix} \quad (1.1)$$

and set $H = \{h(s) : s \in \mathbb{R}\}$.

In the case when the orbit $\Gamma \backslash \Gamma H$ is closed and of finite length, the limiting distribution of the translates $\Gamma \backslash \Gamma H a(T)$ as $T \rightarrow \infty$ is described by the unique G -invariant probability measure $d\mu(g) = dg$ on $\Gamma \backslash G$ [10], that is, if s_0 is the period of $\Gamma \cap H \backslash H$, then for any $\psi \in C_c(\Gamma \backslash G)$,

$$\lim_{T \rightarrow \pm\infty} \frac{1}{s_0} \int_{s=0}^{s_0} \psi(h(s)a(T)) ds = \int_{\Gamma \backslash G} \psi dg.$$

Similarly, understanding the limit of the translates $\Gamma \backslash \Gamma H a(T)$, when $\Gamma \backslash \Gamma H$ is of infinite length, is the main new ingredient in our proofs of Theorem 1.2.

Theorem 1.6. *Let $x_0 \in \Gamma \backslash G$ and suppose that $x_0 h(s)$ diverges as $s \rightarrow +\infty$, that is, $x_0 h(s)$ leaves every compact subset for all sufficiently large $s \gg 1$. For a given compact subset $\Omega \subset \Gamma \backslash G$, there exists $M = M(\Omega, x_0) > 0$ such that for any $\psi \in C^\infty(\Gamma \backslash G)$ with support in Ω , we have, as $|T| \rightarrow \infty$,*

$$\int_0^\infty \psi(x_0 h(s)a(T)) ds = \int_0^{T+M} \psi(x_0 h(s)a(T)) ds = |T| \int \psi d\mu + O(1),$$

where the implied constant depends on x_0 , Ω and $\mathcal{S}^\dagger(\psi) = \max\{\|\psi\|_{C^1}, \mathcal{S}_1(\psi)\}$; here $\|\psi\|_{C^1}$ and $\mathcal{S}_1(\psi)$ respectively mean the C^1 -norm (see (3.3)) and the L^2 -Sobolev norm of ψ of degree one (see (3.1)).

Remark 1.7. Consider the hyperbolic plane \mathbb{H}^2 . A parabolic fixed point for Γ is a point in the geometric boundary $\partial_\infty(\mathbb{H}^2)$ fixed by a parabolic element of Γ . If $\mathcal{F} \subset \mathbb{H}^2$ is a finite sided Dirichlet region for Γ , then the parabolic fixed points of Γ are precisely the Γ -orbits of vertices of $\overline{\mathcal{F}}$ lying in $\partial_\infty(\mathbb{H}^2)$. Let $\pi : G \rightarrow \mathbb{H}^2$ denote the orbit map $g \mapsto g(i)$. For $x_0 = \Gamma g_0 \in \Gamma \backslash G$, the image $\pi(g_0 H)$ is a geodesic in \mathbb{H}^2 with two endpoints $g_0 H(+\infty) := \lim_{s \rightarrow \infty} \pi(g_0 h(s))$ and $g_0 H(-\infty) := \lim_{s \rightarrow -\infty} \pi(g_0 h(s))$ in $\partial_\infty(\mathbb{H}^2)$. We remark that $x_0 h(s)$ diverges as $s \rightarrow +\infty$ (resp. $s \rightarrow -\infty$) if and only if $g_0 H(+\infty)$ (resp. $g_0 H(-\infty)$) is a parabolic fixed point for Γ (cf. Theorem 2.1).

Corollary 1.8. *Suppose that $x_0 H$ is closed and non-compact. For any $\psi \in C_c(\Gamma \backslash G)$,*

$$\lim_{T \rightarrow \pm\infty} \frac{1}{2|T|} \int_{-\infty}^{\infty} \psi(x_0 h(s) a(T)) ds = \int_{\Gamma \backslash G} \psi d\mu.$$

It is well-known due to the work of Duke, Rudnick and Sarnak [10] that Theorem 1.2 follows once we establish the equidistribution of the translates $\Gamma \backslash \Gamma H a(T)$ of a divergent H -orbit as in Theorem 1.6. Our proof of Theorem 1.6 is based on the description of divergent orbits due to Dani [8] and the effective equidistribution of the translates of a small piece of $\Gamma \backslash \Gamma H$ due to Kleinbock and Margulis [15]. The new ingredient of our work is mainly in the section 4.

Acknowledgments We thank Zeev Rudnick for insightful comments.

2. STRUCTURE OF CUSPS IN $\Gamma \backslash G$ AND DIVERGENT TRAJECTORY

Let $G = \mathrm{SL}_2(\mathbb{R})$ and Γ be a non-uniform lattice in G . We will keep the notation for $h(s)$ and $a(s)$ from (1.1) in the introduction. Let

$$N = \left\{ \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} : s \in \mathbb{R} \right\} \quad \text{and} \quad U = w N w^{-1}$$

where $w = \begin{bmatrix} \cos(\pi/4) & \sin(\pi/4) \\ -\sin(\pi/4) & \cos(\pi/4) \end{bmatrix}$. Note that $h(s) = w a(s) w^{-1}$ for all $s \in \mathbb{R}$. For $\eta > 0$, let

$$H_\eta = \{h(s) : s/2 > -\log \eta\}.$$

Let $K = \mathrm{SO}(2) = \{g \in G : g g^t = I\}$. Then the multiplication map $U \times H \times K \rightarrow G : (u, h, k) \mapsto u h k$ is a diffeomorphism.

The following classical result may be found at [13, Thm. 0.6] or [9]:

Theorem 2.1. *There exists a finite set $\Sigma \subset G$ such that the following holds:*

- (1) $\Gamma \backslash \Gamma \sigma U$ is compact for every $\sigma \in \Sigma$.

(2) For any $\eta > 0$, the set

$$\Omega_\eta := \Gamma \backslash G \setminus \bigcup_{\sigma \in \Sigma} \Gamma \backslash \Gamma \sigma U H_\eta K$$

is compact; and any compact subset of $\Gamma \backslash G$ is contained in Ω_η for some $\eta > 0$.

(3) There exists $\eta_0 > 0$ such that for $i = 1, 2$, if $\sigma_i \in \Sigma$, $u_i \in U$, $h_i \in H_{\eta_0}$, and $\Gamma \sigma_1 u_1 h_1 k_1 = \Gamma \sigma_2 u_2 h_2 k_2$, then $\sigma_1 = \sigma_2$, $k_1 = \pm k_2$ and $h_1 = h_2$.

Consider the standard representation of $G = \mathrm{SL}_2(\mathbb{R})$ on \mathbb{R}^2 : $((v_1, v_2), g) \mapsto (v_1, v_2)g$. Let $\|\cdot\|$ denote the Euclidean norm on \mathbb{R}^2 . Let

$$p = (0, 1)w^{-1} = (-\sin(\pi/4), \cos(\pi/4)).$$

Then $pU = p$, and $ph(s) = (0, 1)a(s)w^{-1} = e^{-s/2}p$ for all $s \in \mathbb{R}$. Also

$$g \in UH_\eta K \Leftrightarrow \|pg\| < \eta. \quad (2.1)$$

Proposition 2.2 (Dani [7]). *Let $x_0 \in \Gamma \backslash G$ be such that the trajectory $\{x_0 h(s) : s \geq 0\}$ is divergent. Then there exist $\sigma_0 \in \pm I\Sigma$, $s_0 \in \mathbb{R}$ and $u \in U$ such that $x_0 = \Gamma \sigma_0 u h(s_0)$.*

Proof. By Theorem 2.1, there exists $s_1 > 0$ and $\sigma \in \Sigma$ such that $x_0 h(s) = \Gamma \sigma U H_{\eta_0/2} K$ for all $s \geq s_1$. Let $g_1 \in UH_{\eta_0/2} K$ be such that $x_0 h(s_1) = \Gamma \sigma g_1$. We claim that $pg_1 \in \mathbb{R}p$. If not, then $\|pg_1 h(s)\| \rightarrow \infty$ as $s \rightarrow \infty$, and hence there exists $s > 0$ such that $\eta_0/2 < \|pg_1 h(s)\| < \eta_0$. By (2.1), $g_1 h(s) = uhk$ for some $u \in U$, $h \in H_{\eta_0}$ and $k \in K$. Therefore

$$\Gamma \sigma u h k = \Gamma \sigma g_1 h(s) = x_0 h(s_1 + s) \in \Gamma \sigma U H_{\eta_0/2} K.$$

By Theorem 2.1(iii), we have that $h \in H_{\eta_0/2}$. But then $\|pg_1 h(s)\| = \|puhk\| < \eta_0/2$, a contradiction. Therefore our claim that $pg_1 \in \mathbb{R}p$ is valid. Hence $g_1 = u_1 h(s) \{\pm I\}$ for some $u_1 \in U$ and $s/2 \geq -\log(\eta_0/2)$. Thus $x_0 h(s_1) = \Gamma \sigma u_1 h(s) \{\pm I\}$, and hence $x_0 = \Gamma \sigma_0 u_1 h(s - s_1)$, where $\sigma_0 = \pm I\sigma$. \square

Proposition 2.3. *Let $x_0 \in \Gamma \backslash G$ be such that the trajectory $\{x_0 h(s) : s \geq 0\}$ is divergent. Let $\Omega \subset \Gamma \backslash G$ be a compact subset. There exists $M_1 = M_1(\Omega, x_0) > 0$ such that*

$$x_0 h(s) a(T) \notin \Omega$$

for any $T \in \mathbb{R}$ and $s > 0$ satisfying $s > |T| + M_1$. In particular, for any $f \in C(\Gamma \backslash G)$ with support inside Ω ,

$$\int_0^\infty f(x_0 h(s) a(T)) ds = \int_0^{|T|+M_1} f(x_0 h(s) a(T)) ds.$$

Proof. By Proposition 2.2, $x_0 = \Gamma \sigma_0 u h(s_0)$ for some $\sigma_0 \in \pm \Sigma$, $u \in U$, $s_0 \in \mathbb{R}$. By Theorem 2.1(ii), let $\eta > 0$ be such that $\Omega \subset \Omega_\eta$. Let $M_1 = -s_0 - 2 \log(\eta)$.

Since $s - |T| > -s_0 - 2 \log \eta$, we have

$$\begin{aligned} \|puh(s_0)h(s)a(T)\| &= \|ph(s+s_0)a(T)\| \\ &= e^{-(s+s_0)/2} \|pa(T)\| \\ &< e^{-(s+s_0)/2} e^{|T|/2} \\ &= e^{-(s+s_0-|T|)/2} < \eta. \end{aligned} \quad (2.2)$$

Therefore by (2.1), $uh(s_0)h(s)a(T) \in UH_\eta K$, and hence

$$x_0h(s)a(T) \in \Gamma\sigma_0 UH_\eta K \subset \Gamma \backslash G \setminus \Omega_\eta.$$

□

3. UNIFORM MIXING ON COMPACT SETS

Let $G = \mathrm{SL}_2(\mathbb{R})$ and $\Gamma < G$ be a lattice. Let μ denote the G -invariant probability measure on $\Gamma \backslash G$. For an orthonormal basis X_1, X_2, X_3 of $\mathfrak{sl}(2, \mathbb{R})$ with respect to an Ad -invariant scalar product, and $\psi \in C^\infty(\Gamma \backslash G)$, we consider the Sobolev norm

$$\mathcal{S}_m(\psi) = \max\{\|X_{i_1} \cdots X_{i_j}(\psi)\|_2 : 1 \leq i_j \leq 3, 0 \leq j \leq m\}. \quad (3.1)$$

The well-known spectral gap property for $L^2(\Gamma \backslash G)$ says that the trivial representation is isolated (see [1, Lemma 3]) in the Fell topology of the unitary dual of G . It follows that there exist $\theta > 0$ and $c > 0$ such that for any $\psi_1, \psi_2 \in C^\infty(\Gamma \backslash G)$ with $\int \psi_i d\mu = 0$, $\mathcal{S}_1(\psi_i) < \infty$ and for any $T > 0$,

$$|\langle a(T)\psi_1, \psi_2 \rangle| \leq ce^{-\theta|T|} \mathcal{S}_1(\psi_1) \mathcal{S}_1(\psi_2) \quad (3.2)$$

(cf. [5], [21])

Write $\mathcal{O}_\epsilon = \{g \in G : \|g - I\|_\infty \leq \epsilon\}$. For a compact subset $\Omega \subset \Gamma \backslash G$, let $0 < \epsilon_0(\Omega) \leq 1$ be the injectivity radius of Ω , that is, $\epsilon_0(\Omega)$ is the supremum of $0 < \epsilon \leq 1$ such that the multiplication map $\Omega \times \mathcal{O}_\epsilon \rightarrow \Gamma \backslash G$ is injective.

For $s \in \mathbb{R}$, let

$$n_+(s) = \begin{pmatrix} 1 & 0 \\ s & 1 \end{pmatrix} \quad \text{and} \quad n_-(s) = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix}.$$

Let $\|f\|_{C^1}$ denotes the C^1 -norm of f , that is,

$$\|f\|_{C^1} = \|f\|_\infty + \sum_{i=1}^3 \|X_i(f)\|_\infty. \quad (3.3)$$

Set $\mathcal{S}^\dagger(f) := \max\{\mathcal{S}_1(f), \|f\|_{C^1}\}$; in fact, $\mathcal{S}_1(f) \leq \|f\|_{C^1}$.

The following is a special case of [15, Prop. 2.4.8]:

Theorem 3.1. *Let $\Omega \subset \Gamma \backslash G$ be a compact subset and $\eta > 0$. There exists $c = c(\Omega) > 0$ such that for any $\psi \in C^\infty(\Gamma \backslash G)$ with support in Ω , for any $|T| \geq 1$, $x \in \Omega$, and $0 < r_0 < \epsilon_0(\Omega)$, we have*

$$\left| \int_0^{r_0} \psi(xn_\nu(s)a(T)) ds - r_0 \int \psi d\mu \right| \leq c(\mathcal{S}^\dagger(\psi) + 1)e^{-\theta_0|T|} \quad (3.4)$$

for some θ_0 depending only on the spectral gap for $L^2(\Gamma \backslash G)$. Here and in what follows, the sign $\nu = +$ if $T > 0$ and $\nu = -$ if $T < 0$.

4. TRANSLATES OF DIVERGENT ORBITS

Let $x_0 \in \Gamma \backslash G$ be such that $x_0 h(s)$ diverge as $s \rightarrow \infty$.

Theorem 4.1. *For any $|T| > 1$ and any $\psi \in C_c^\infty(\Gamma \backslash G)$*

$$\int_0^{|T|} \psi(x_0 h(s) a(T)) ds = |T| \int \psi d\mu + O(1) \mathcal{S}^\dagger(\psi)$$

where the implied constant depends on x_0 , $\mathcal{S}^\dagger(\psi)$ and the support of ψ .

Proof. Let $R_0 = -\log \eta_0$. Due to Proposition 2.2, replacing x_0 by another point in $x_0 H$, we may assume that $x_0 = \Gamma \sigma_0 h(R_0)$. For any $S > 0$, $\|ph(R_0)h(S)a(S)\| \in [\eta_0/\sqrt{2}, \eta_0]$. Hence $x_0 h(R_0)h(S)a(S) \in \Omega_{\eta_0/\sqrt{2}}$.

Let r_0 be the injectivity radius of $\Omega_{\eta_0/\sqrt{2}}$, that is, $r_0 = \epsilon_0(\Omega_{\eta_0/\sqrt{2}})$. Let $S_0 = 0$, and choose S_i such that $r_0 e^{-S_i} \leq \delta_i := S_{i+1} - S_i \leq 2r_0 e^{-S_i}$ for each i . We will choose $S_i = \log(2r_0 i + 1)$ for each i . Then $x_0 h(S_i) a(S_i) \in \Omega_{\eta_0/\sqrt{2}}$. We put $R_i = T - S_i$.

We will express $x_0 h([S_i, S_{i+1}]) a(T) = x_i h^{a(S_i)}([0, \delta_i]) a(R_i)$, where $x_i = x_0 h(S_i) a(S_i)$, $h^{a(S_i)}(s) = a(-S_i) h(s) a(S_i) = n(e^{S_i} s/2) w_i(s)$, and $|w_i(s)| = O(e^{-2S_i})$. Note that $r_0/2 \leq e^{S_i} \delta_i/2 \leq r_0$.

By Theorem 3.1, we have

$$\int_0^{r_0} \psi(x_i n(s) a(R_i)) ds - r_0 \int \psi d\mu = \mathcal{S}^\dagger(\psi) \cdot O(e^{-\theta_0 R_i})$$

and hence

$$\int_{S_i}^{S_{i+1}} \psi(x_0 h(s) a(T)) ds = \frac{\delta_i}{r_0} \int_0^{r_0} \psi(x_i n(s) a(R_i)) ds + \mathcal{S}^\dagger(\psi) \cdot O(e^{-2S_i} \delta_i).$$

Let $k = k(T)$ be such that $S_k \leq T < S_k + r_0 e^{-S_k}$. Therefore, since $\delta_i r_0^{-1} \leq 2e^{-S_i}$,

$$\begin{aligned} \int_0^T \psi(xh(s)a(T)) ds &= \sum_{i=0}^{k-1} \int_{S_i}^{S_{i+1}} \psi(xh(s)a(T)) ds + O(e^{-S_k}) \\ &= \sum_{i=0}^{k-1} \delta_i \frac{1}{r_0} \int_0^{r_0} \psi(x_i n(s) a(T)) ds + \mathcal{S}^\dagger(\psi) \cdot O(e^{-2S_i} \delta_i) + O(1) \\ &= \sum_{i=0}^{k-1} \delta_i \mu(\psi) + \sum_{i=0}^{k-1} \delta_i r_0^{-1} \mathcal{S}^\dagger(\psi) \cdot O(e^{-\theta_0 R_i}) + \mathcal{S}^\dagger(\psi) \cdot O(e^{-2S_i} \delta_i) + O(1) \\ &= T\mu(\psi) + O\left(\sum_{i=1}^{k-1} e^{-S_i} e^{-\theta_0 R_i} + \sum_{i=1}^k e^{-3S_i}\right) \mathcal{S}^\dagger(\psi) + O(1) \\ &= T\mu(\psi) + O\left(e^{-\theta_0 T} \sum_{i=0}^{k-1} e^{(1-\theta_0)S_i} + \sum_{i=0}^{k-1} e^{-3S_i}\right) \mathcal{S}^\dagger(\psi) + O(1). \end{aligned}$$

Since $S_i = \log(2r_0i + 1)$, $0 < T - S_k < 2e^{-T}$ implies that $k < \frac{e^T - 1}{2r_0} < k + 1$, and hence

$$\sum_{i=0}^{k-1} e^{-3S_i} \ll \sum_{i=1}^{k-1} \frac{1}{(2r_0i + 1)^3} = O(k^{-2} + 1) = O(e^{-2T} + 1) < \infty$$

and

$$\sum_{i=0}^{k-1} e^{(1-\theta_0)S_i} \ll \int_0^{e^T} \frac{1}{(2r_0x + 1)^{1-\theta_0}} dx = O(e^{\theta_0 T}).$$

Hence

$$e^{-\theta_0 T} \sum_{i=0}^{k-1} e^{(1-\theta_0)S_i} + \sum_{i=0}^{k-1} e^{-3S_i} = O(1).$$

Therefore

$$\int_0^T \psi(xh(s)a(T)) ds = T\mu(\psi) + O(1)\mathcal{S}^\dagger(\psi).$$

□

Theorem 1.6 follows from the following:

Theorem 4.2. *Let $x_0h(s)$ diverge as $s \rightarrow \infty$. For a given compact subset $\Omega \subset \Gamma \backslash G$, and $\psi \in C^\infty(\Gamma \backslash G)$ with support in Ω , we have*

$$\int_0^\infty \psi(x_0h(s)a(T)) ds = |T| \cdot \int \psi d\mu + O(1)\mathcal{S}^\dagger(\psi)$$

with the implied constant depending only on x_0 , $\mathcal{S}^\dagger(\psi)$, and the support of ψ .

Proof. Since $x_0h(s)$ diverges as $s \rightarrow \infty$, by Proposition 2.3, there exists $M_1 = M_1(\Omega) > 0$ such that

$$\begin{aligned} \int_0^\infty \psi(x_0h(s)a(T)) ds &= \int_0^{|T|+M_1} \psi(x_0h(s)a(T)) ds \\ &= (|T| + M_1) \int \psi d\mu + O(1)\mathcal{S}^\dagger(\psi) \\ &= |T| \int \psi d\mu + O(1)\mathcal{S}^\dagger(\psi). \end{aligned}$$

□

By a similar argument, we also deduce the following:

Corollary 4.3. *If $x_0h(s)$ diverges as $s \rightarrow -\infty$, then*

$$\int_{-\infty}^0 \psi(x_0h(s)a(T)) ds = |T| \int \psi d\mu + O(1)\mathcal{S}^\dagger(\psi)$$

with the implied constant depending only on x_0 , $\mathcal{S}^\dagger(\psi)$, and the support of ψ .

Lemma 4.4. *If $x_0h(\mathbb{R})$ is closed and non-compact, then $x_0h(s)$ diverges as $s \rightarrow \pm\infty$.*

Proof. We use a well-known fact that for a closed subgroup H of a locally compact second countable group G and a discrete subgroup Γ of G , if ΓH is closed in G , then the canonical projection map $H \cap \Gamma \backslash H \rightarrow \Gamma \backslash G$ is a proper map (cf. [18]). Since $x_0h(\mathbb{R})$ is non-compact and $h(\mathbb{R})$ is one-dimensional with no non-trivial finite subgroups, the stabilizer of x_0 in $h(\mathbb{R})$ is trivial. Therefore the map $h(\mathbb{R}) \rightarrow \Gamma \backslash G$ given by $h \rightarrow x_0h$ is a proper injective map. This implies that $x_0h(s)$ diverges as $s \rightarrow \pm\infty$. \square

Proof of Corollary 1.8. As the set $C_c^\infty(\Gamma \backslash G)$ is dense in $C_c(\Gamma \backslash G)$, the claim follows from Lemma 4.4, Theorem 4.1, and Corollary 4.3. \square

5. COUNTING: PROOF OF THEOREM 1.2

Let Q be a real quadratic form in 3 variables of signature $(2, 1)$ and Γ_0 a lattice in the identity component G_0 of $\mathrm{SO}_Q(\mathbb{R})$. We assume that $v_0\Gamma_0$ is discrete for some vector $v_0 \in \mathbb{R}^3$ with $Q(v_0) = d > 0$ and that the stabilizer H_0 of v_0 in G_0 is finite.

It suffices to prove Theorem 1.2 in the case when $Q = x^2 + y^2 - z^2$ and $v_0 = (\sqrt{d}, 0, 0)$ by the virtue of Witt's theorem.

Consider the spin double cover map $\iota : G := \mathrm{SL}_2(\mathbb{R}) \rightarrow G_0$ given by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} \frac{a^2-b^2-c^2+d^2}{2} & ac-bd & \frac{a^2-b^2+c^2-d^2}{2} \\ ab-cd & bc+ad & ab+cd \\ \frac{a^2+b^2-c^2-d^2}{2} & ac+bd & \frac{a^2+b^2+c^2+d^2}{2} \end{pmatrix}.$$

For $s \in \mathbb{R}$ and $\theta \in [0, 2\pi)$, we set

$$h(s) = \begin{pmatrix} \cosh(s/2) & \sinh(s/2) \\ \sinh(s/2) & \cosh(s/2) \end{pmatrix}; \quad a(s) = \begin{pmatrix} e^{s/2} & 0 \\ 0 & e^{-s/2} \end{pmatrix} \quad \text{and} \quad k(\theta) = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}.$$

Recall that $H := \{h(s) : s \in \mathbb{R}\}$, $A := \{a(t) : t \in \mathbb{R}\}$ and $K_1 := \{k(\theta) : \theta \in [0, 2\pi)\}$, here K_1 is half of the circle group. Observing that

$$\iota(h(s)) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cosh s & \sinh s \\ 0 & \sinh s & \cosh s \end{pmatrix} \quad \text{and} \quad \iota(a(t)) = \begin{pmatrix} \cosh t & 0 & \sinh t \\ 0 & 1 & 0 \\ \sinh t & 0 & \cosh t \end{pmatrix},$$

the subgroup $\tilde{H} := \pm H$ is the stabilizer of v_0 in G . We have a generalized Cartan decomposition $G = \tilde{H}AK_1$ in the sense that every g is of the form hak for unique $h \in \tilde{H}, a \in A, k \in K_1$. And for $g = h(s)a(t)k$, $d\mu(g) = \sinh(t)dsdtdk$ defines a Haar measure on G , where $dk = (1/2\pi)dk(\theta)$, and ds, dt and $d\theta$ are Lebesgue measures. As $v_0G = \pm H \backslash G \simeq A \times K_1$, $\sinh(t)dtdk$ defines an invariant measure on v_0G . We consider the volume forms on G and v_0G with respect to these measures. Via the map ι , these define invariant measures on G_0 and v_0G_0 as well.

Denote by Γ the pre-image of Γ_0 under ι . Then $\mathrm{Stab}_\Gamma(v_0) = \tilde{H} \cap \Gamma = \{\pm I\}$.

For each $T > 1$, define a function on $\Gamma \backslash G$:

$$F_T(g) := \sum_{\gamma \in \pm I \backslash \Gamma} \chi_{B_T}(v_0 \gamma g).$$

Proposition 5.1. *For any $\Psi \in C_c^\infty(\Gamma \backslash G)$,*

$$\langle F_T, \Psi \rangle = \frac{T \log T \mu(\Psi)}{\text{vol}(\Gamma \backslash G)} \cdot 2 \int_{K_1} \frac{1}{\|v^+ k\|} dk + O(\mathcal{S}^\dagger(\Psi)T)$$

where $v^\pm = \frac{\sqrt{d}}{2}(e_1 \pm e_3)$. Here the implied constant depends only on the support of Ψ and $\mathcal{S}^\dagger(\Psi)$.

Proof. Note that $v_0 = v^+ + v^-$ and $v_0 a(t) = e^t v^+ + e^{-t} v^-$. Since $B_T = \{v_0 a(t)k : \|v_0 a(t)k\| < T, t \in \mathbb{R}, k \in K_1\}$, we have

$$\begin{aligned} \langle F_T, \Psi \rangle &= \int_{\Gamma \backslash G} \sum_{\gamma \in \pm I \backslash \Gamma} \chi_{B_T}(v_0 \gamma g) \Psi(g) d\mu(g) \\ &= \int_{k \in K_1} \int_{\|v_0 a(t)k\| < T} \left(\int_{h(s) \in \pm I \backslash \tilde{H}} \Psi(h(s)a(t)k) ds \right) \sinh(t) dt dk \\ &= \int_{k \in K_1} \int_{\|v_0 a(t)k\| < T} \left(\int_{s \in \mathbb{R}} \Psi(h(s)a(t)k) ds \right) \sinh(t) dt dk. \end{aligned}$$

Since $v_0 \Gamma$ is discrete and $H \cap \Gamma$ is trivial, it follows that $\Gamma \backslash \Gamma H$ is closed and non-compact in $\Gamma \backslash G$. Now fix any $k \in K_1$. Hence by Theorem 4.2 and Lemma 4.4,

$$\begin{aligned} &\int_{t \gg 1, \|v_0 a(t)k\| < T} \left(\int_{s \in \mathbb{R}} \Psi(h(s)a(t)k) ds \right) \sinh(t) dt \\ &= \frac{1}{\text{vol}(\Gamma \backslash G)} \int_{t \gg 1, e^t \|v^+ k\| < T + O(1)} (2t\mu(\Psi) + O(1)\mathcal{S}^\dagger(\Psi))(e^t/2 + O(1)) dt \\ &= \frac{T \log T \mu(\Psi)}{\text{vol}(\Gamma \backslash G) \cdot \|v^+ k\|} + O(T)\mathcal{S}^\dagger(\Psi). \end{aligned}$$

Similarly,

$$\begin{aligned} &\int_{t \ll -1, \|v_0 a(t)k\| < T} \left(\int_{s \in \mathbb{R}} \Psi(h(s)a(t)k) ds \right) \sinh(t) dt \\ &= \int_{t \gg 1, \|v_0 a(-t)k\| < T} \left(\int_{s \in \mathbb{R}} \Psi(h(s)a(-t)k) ds \right) \sinh(t) dt \\ &= \frac{1}{\text{vol}(\Gamma \backslash G)} \int_{t \gg 1, e^t \|v^- k\| < T + O(1)} (2t\mu(\Psi) + O(1)\mathcal{S}^\dagger(\Psi))(e^t/2 + O(1)) dt \\ &= \frac{T \log T \mu(\Psi)}{\text{vol}(\Gamma \backslash G) \|v^- k\|} + O(T)\mathcal{S}^\dagger(\Psi). \end{aligned}$$

Since $v^-k(\pi) = -v^+$,

$$\int_{k \in K_1} \|v^-k\|^{-1} dk = \int_{k \in K_1} \|v^+k(\pi)k\|^{-1} dk = \int_{K_1} \|v^+k\|^{-1} dk.$$

The required formula can be deduced in a straightforward manner from this. \square

Fix a non-negative function $\psi \in C_c^\infty(G)$ whose support injects to $\Gamma \backslash G$ and with integral $\int \psi(g) d\mu(g) = 1$. Consider a function ξ_T on \mathbb{R}^3 defined by

$$\xi_T(x) = \int_{g \in G} \chi_{B_T}(xg) \psi(g) d\mu(g).$$

Then the sum $\sum_{\gamma \in \pm I \backslash \Gamma} \xi_T(v_0\gamma)$ is a smoothed over counting satisfying

$$\sum_{\gamma \in \pm I \backslash \Gamma} \xi_T(v_0\gamma) \asymp \#v_0\Gamma \cap B_T.$$

Theorem 5.2. *As $T \rightarrow \infty$,*

$$\sum_{\gamma \in \pm I \backslash \Gamma} \xi_T(v_0\gamma) = \frac{2T \log T}{\text{vol}(\Gamma \backslash G)} \cdot \int_{k \in K_1} \frac{1}{\|v^+k\|} dk + O(T).$$

Proof. It is not hard to verify that

$$\sum_{\gamma \in \pm I \backslash \Gamma} \xi_T(v_0\gamma) = \langle F_T, \Psi \rangle$$

where $\Psi(\Gamma g) = \sum_{\gamma \in \Gamma} \psi(\gamma g)$. Therefore the claim follows from Proposition 5.1. \square

Theorem 5.3. *For $T \gg 1$, we have*

$$\#\{w \in v_0\Gamma : \|w\| < T\} = \frac{2T \log T}{\text{vol}(\Gamma \backslash G)} \int_{K_1} \frac{1}{\|v^+k\|} dk \cdot (1 + (\log T)^{-\alpha})$$

where $\alpha = 0.25$.

Proof. Note that $F_T(I) = \#\{w \in v_0\Gamma : \|w\| < T\}$. For each $\epsilon > 0$, let $\mathcal{O}_\epsilon = \{g \in G : \|g - I\|_\infty \leq \epsilon\}$. There exists $0 < \ell \leq 1$ such that for all small $\epsilon > 0$,

$$\mathcal{O}_{\ell\epsilon} B_T \subset B_{(1+\epsilon)T}, \quad B_{(1-\epsilon)T} \subset \cap_{u \in \mathcal{O}_{\ell\epsilon}} u B_T. \quad (5.1)$$

Let ψ^ϵ be a non-negative smooth function on G supported in $\mathcal{O}_{\ell\epsilon}$ and with integral $\int \psi^\epsilon d\mu = 1$ and define $\Psi^\epsilon \in C_c^\infty(\Gamma \backslash G)$ by $\Psi^\epsilon(\Gamma g) := \sum_{\gamma \in \Gamma} \psi^\epsilon(\gamma g)$.

Using (5.1), we have

$$\langle F_{(1-\epsilon)T}, \Psi^\epsilon \rangle \leq F_T(I) \leq \langle F_{(1+\epsilon)T}, \Psi^\epsilon \rangle.$$

Note that $\mathcal{S}_1(\Psi^\epsilon) = O(\epsilon^{-3/2})$ and $\|(\Psi^\epsilon)\|_{C^1} = O(\epsilon^{-3})$ so that $\mathcal{S}^\dagger = O(\epsilon^{-3})$. Therefore by Proposition 5.1

$$\begin{aligned} \langle F_{(1\pm\epsilon)T}, \Psi^\epsilon \rangle &= \frac{2T \log T}{\text{vol}(\Gamma \backslash G)} \int_{K_1} \frac{1}{\|v+k\|} dk + O(\epsilon T \log T) + O(\mathcal{S}^\dagger(\Psi^\epsilon)T) \\ &= \frac{2T \log T}{\text{vol}(\Gamma \backslash G)} \int_{K_1} \frac{1}{\|v+k\|} dk + O(\epsilon T \log T) + O(\epsilon^{-3}T) \\ &= \frac{2T \log T}{\text{vol}(\Gamma \backslash G)} \int_{K_1} \frac{1}{\|v+k\|} dk (1 + (\log T)^{-1/4}), \end{aligned}$$

where the last equality follows by we putting $\epsilon = (\log T)^{-1/4}$. \square

Proof of Theorem 1.2. The computation in the proof of Proposition 5.1 also shows that

$$\text{vol}(B_T) = \int_{k \in K_1} \int_{\|v_0 a(t)k\| < T} \sinh(t) dt dk = T \int_{k \in K} \frac{1}{\|v+k\|} dk + O(\log T). \quad (5.2)$$

From Theorem 5.3, it follows that

$$F_T(I) = \frac{2 \log T \text{vol}(B_T)}{\text{vol}(\Gamma \backslash G)} (1 + O(\log T)^{-\alpha}). \quad (5.3)$$

Since $F_T(I) = \#(v_0 \Gamma \cap B_T)$, this completes the proof of the first claim (1). The second claim (2) follows from Proposition 5.1. \square

6. ORBITAL COUNTING FOR GENERAL REPRESENTATIONS OF $\text{SL}_2(\mathbb{R})$

Let $G = \text{SL}_2(\mathbb{R})$ and Γ be a non-uniform lattice in G . For $s \in \mathbb{R}$, define

$$h(s) = \begin{bmatrix} \cosh(s/2) & \sinh(s/2) \\ \sinh(s/2) & \cosh(s/2) \end{bmatrix}, \quad a(s) = \begin{bmatrix} e^{s/2} & 0 \\ 0 & e^{-s/2} \end{bmatrix}, \quad k(\theta) = \begin{bmatrix} \cos(\theta/2) & -\sin(\theta/2) \\ \sin(\theta/2) & \cos(\theta/2) \end{bmatrix}$$

Put $H = \{h(s) : s \in \mathbb{R}\}$, $A^+ = \{a(t) : t > 0\}$, and $K_1 = \{k(\theta) : \theta \in [0, 2\pi]\}$, here K_1 is half of the circle group. Put $w_0 = k(\pi)$. Then $\{\pm I\} \backslash G = HA^+K_1 \cup Hw_0A^+K_1$, $w_0^{-1}h(s)w_0 = h(-s)$ and $w_0^{-1}a(t)w_0 = a(-t)$.

Let V be any finite dimensional representation of G and $v_0 \in G$ be such that H is the stabilizer subgroup of v_0 in G , i.e., $H = G_{v_0}$ where $G_{v_0} = \{g \in G : v_0 g = v_0\}$. Assume that V is linearly spanned by $v_0 G$. Then if e^{mt} is the highest eigenvalue for $a(t)$ -action on V , then $m \in \mathbb{N}$, and the G action factors through $\{\pm I\} \backslash G = \text{PSL}_2(\mathbb{R}) \cong \text{SO}(2, 1)^0$.

For example, let V_m denote the $(2m+1)$ -dimensional space of real homogeneous polynomials of degree $2m$ in two variables, and consider the standard right action of $g \in \text{SL}(2, \mathbb{R})$ on $P(x, y) \in V_m$ by $(Pg)(x, y) = P((x, y)g)$, where $(x, y) \begin{bmatrix} a & b \\ c & d \end{bmatrix} = (ax + cy, bx + dy)$. Let $v_0(x, y) = (x^2 - y^2)^m$. Then $G_{v_0} = H\mathcal{W}$, where $\mathcal{W} = \{\pm I\}$ if m is odd and $\mathcal{W} = \{\pm I, \pm w_0\}$ if m is even. Moreover, $\{P \in V_m : Ph = P, \text{ for all } h \in H\} = \mathbb{R}P_0$. A general finite dimensional representation of G with a nonzero H -fixed vector is a direct sum of such irreducible representations, and v_0 is a sum of one nonzero H -fixed

vector from each of the irreducible representations; we assume that V is a span of v_0G .

Theorem 6.1. *Let V , v_0 and m be as above. Suppose that Γ is a lattice in G , $v_0\Gamma$ is discrete, and $\Gamma_{v_0} := \Gamma \cap G_{v_0}$ is finite. Let $\|\cdot\|$ be any norm on V , and $v_0^+ = \lim_{t \rightarrow \infty} v_0 a_t / \|v_0 a_t\|$. Let C be an open subset of $\{v \in V : \|v\| = 1\}$ such that $\Theta = \{\theta \in [0, 2\pi] : v_0^+ k(\theta) \in \mathbb{R}C\}$ has positive Lebesgue measure, and $\{\theta \in [0, 2\pi] : v_0^+ k(\theta) \in \mathbb{R}(\overline{C} \setminus C)\}$ has zero Lebesgue measure. Then for $T \gg 1$,*

$$\begin{aligned} \#(v_0\Gamma \cap [0, T]C) & \qquad \qquad \qquad (6.1) \\ &= \frac{4(2\pi)^{-1} \int_{\Theta} \|v_0^+ k(\theta)\|^{-1/m} d\theta}{|\Gamma_{v_0}| \cdot \text{vol}_G(\Gamma \backslash G)} \times \frac{\log T}{m} T^{1/m} (1 + (\log T)^{-\alpha}) \end{aligned}$$

where vol_G is given by the Haar integral $dg = \sinh(t) dt ds d\theta$ on G , where $g = h(s)a(t)k(\theta)$, and $\alpha = \frac{1}{4}$.

Moreover, if $C \subset V$ satisfies $\mathbb{R}\overline{C} \cap v_0^+ K_1 = \emptyset$, then $\#(v_0\Gamma \cap \mathbb{R}C) < \infty$.

Proof. The result can be deduced by the arguments as in the proof of Theorem 5.3; one may also use the basic ideas from [14] about using the highest weight. \square

REFERENCES

- [1] M. B. Bekka. On uniqueness of invariant means. *Proc. Amer. Math. Soc.*, 126(2):507–514, 1998.
- [2] Yves Benoist and Hee Oh. Effective equidistribution of S-integral points on symmetric varieties. *To appear in Annales de L'Institut Fourier*, arXiv:0706.1621
- [3] A. Borel and Harish-Chandra. Arithmetic subgroups of algebraic groups, *Ann. of Math. (2)*, vol 75, 485–535, 1962.
- [4] J. Buchmann and U. Vollmer. Binary quadratic forms: An algorithmic Approach, *Springer-Verlag, Algorithms and Computation in Mathematics*, Vol 20.
- [5] M. Cowling, U. Haagerup, and R. Howe. Almost L^2 matrix coefficients. *J. Reine Angew. Math.*, 387:97–110, 1988.
- [6] S. G. Dani. On invariant measures, minimal sets and a lemma of Margulis. *Invent. Math.*, 51(3):239–260, 1979.
- [7] S. G. Dani. Divergent trajectories of flows on homogeneous spaces and Diophantine approximation. *J. Reine Angew. Math.*, 359:55–89, 1985.
- [8] S. G. Dani. Orbits of horospherical flows. *Duke Math. J.*, 53(1):177–188, 1986.
- [9] S. G. Dani and J. Smillie. Uniform distribution of horocycle orbits for Fuchsian groups. *Duke Math. J.*, 51(1):185–194, 1984.
- [10] W. Duke, Z. Rudnick, and P. Sarnak. Density of integer points on affine homogeneous varieties. *Duke Math. J.*, 71(1):143–179, 1993.
- [11] A. Eskin and C.T. McMullen. Mixing, counting, and equidistribution in Lie groups. *Duke Math. J.*, 71(1):181–209, 1993.
- [12] A. Eskin, S. Mozes, and N. Shah. Unipotent flows and counting lattice points on homogeneous varieties. *Ann. of Math. (2)*, 143(2):253–299, 1996.
- [13] H. Garland and M. S. Raghunathan. Fundamental domains for lattices in R-rank 1 semisimple Lie groups. *Ann. of Math. (2)*, 92:279–326, 1970.
- [14] A. Gorodnik, H. Oh, and N. Shah. Integral points on symmetric varieties and Satake compactifications. *Amer. J. Math.*, 131(1):1–57, 2009.

- [15] D. Kleinbock and G.A. Margulis. Bounded orbits of Nonquasiunipotent flows on homogeneous spaces In *Amer. Math. Soc. Transl. (2)*, 171, pages 141–172, 1996.
- [16] G.A. Margulis. Dynamical and ergodic properties of subgroup actions on homogeneous spaces with applications to number theory. In *Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990)*, pages 193–215, Tokyo, 1991. Math. Soc. Japan.
- [17] M.Ratner. On Raghunathan’s measure conjecture. *Ann. of Math. (2)*, 134(3):545–607, 1991.
- [18] Hee Oh and Nimish Shah. Equidistribution and counting for orbits of geometrically finite hyperbolic groups. *arxiv:1001.2096*, To appear in J. Amer. Math. Soc.
- [19] E.J. Scourfield. The divisors of a quadratic polynomial. *Proc. Glasgow Math. Assoc.*, 5:8–20, 1961.
- [20] N.A. Shah. Uniformly distributed orbits of certain flows on homogeneous spaces. *Math. Ann.*, 289:315–334, 1991.
- [21] A. Venkatesh. Sparse equidistribution problem, period bounds, and subconvexity, 2007. *Ann.of Math.*, 172:989-1094, 2010.

E-mail address: `heeoh@math.brown.edu`

MATHEMATICS DEPARTMENT, BROWN UNIVERSITY, PROVIDENCE, RI, U.S.A AND KOREA INSTITUTE FOR ADVANCED STUDY, SEOUL, KOREA

E-mail address: `shah@math.osu.edu`

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS, OH 43210, U.S.A